

Convergent dopamine and ALK4 signaling to PCBP1 controls FosB alternative splicing and cocaine behavioral sensitization

Favio A Krapacher¹ , Diana Fernández-Suárez¹, Annika Andersson¹, Alvaro Carrier-Ruiz¹  & Carlos F Ibáñez^{1,2,3,*} 

Abstract

Δ FosB is an alternatively spliced product of the FosB gene that is essential for dopamine-induced reward pathways and that acts as a master switch for addiction. However, the molecular mechanisms of its generation and regulation by dopamine signaling are unknown. Here, we report that dopamine D1 receptor signaling synergizes with the activin/ALK4/Smad3 pathway to potentiate the generation of Δ FosB mRNA in medium spiny neurons (MSNs) of the nucleus accumbens (NAc) *via* activation of the RNA-binding protein PCBP1, a regulator of mRNA splicing. Concurrent activation of PCBP1 and Smad3 by D1 and ALK4 signaling induced their interaction, nuclear translocation, and binding to sequences in exon-4 and intron-4 of FosB mRNA. Ablation of either ALK4 or PCBP1 in MSNs impaired Δ FosB mRNA induction and nuclear translocation of Δ FosB protein in response to repeated co-stimulation of D1 and ALK4 receptors. Finally, ALK4 is required in NAc MSNs of adult mice for behavioral sensitization to cocaine. These findings uncover an unexpected mechanism for Δ FosB generation and drug-induced sensitization through convergent dopamine and ALK4 signaling.

Keywords activin; addiction; drugs of abuse; Δ FosB

Subject Categories Molecular Biology of Disease; Neuroscience

DOI 10.15252/emj.2022110721 | Received 19 January 2022 | Accepted 25 May 2022

The EMBO Journal (2022) e110721

Introduction

The central dopamine system of the mammalian brain, notably the ventral tegmental area (VTA), substantia nigra pars compacta (SNc), and their target neurons in the nucleus accumbens (NAc), has a pivotal role in the regulation of responses to natural rewards such as food, drink, sex and social interaction (Volkow & Morales, 2015; Kauer & Malenka, 2007). One of its main functions is to reinforce behaviors that lead to successful earning of such

rewards to the extent they enhance survival and reproduction. On the flip side, this also makes the dopamine system a vulnerable target that can be hijacked by a variety of psychoactive stimuli, including drugs of abuse, and compulsive behaviors, such as pathological overeating, compulsive running, and gambling (Lüscher, 2016). An important area of neuroscience research is centered on understanding the intricate ways in which drugs of abuse alter the brain to produce the functional and behavioral abnormalities that characterize addiction. The persistence of these alterations is thought to be sustained by transcriptional changes that maintain long-lasting molecular and structural adaptations in neuronal circuits involved in reward processing and its behavioral manifestation (Teague & Nestler, 2021). A key component of transcriptional mechanisms underlying reward responses is Δ FosB, a transcription factor encoded by a splice variant of FosB, a member of the Fos family of immediate early genes (Ruffle, 2014; Nestler, 2013, 2008). Fos family proteins are rapidly induced in the NAc after acute exposure to many drugs of abuse, such as cocaine, but their levels return to baseline within hours of drug administration due to protein instability (Hope *et al.*, 1992; Young *et al.*, 1991; Graybiel *et al.*, 1990). Upon chronic drug exposure, most Fos family proteins show desensitization and fail to accumulate to significant levels (Alibhai *et al.*, 2007). In contrast, repeated drug administration results in the accumulation of Δ FosB, which, due to its unusually high stability, can persist for days or weeks in NAc neurons (Chen *et al.*, 1997, 1995; Hiroi *et al.*, 1997). Δ FosB mRNA is generated by the removal of intron 4 (also known as the regulated intron) from the 4th and last exon of FosB mRNA, thereby introducing a frameshift that leads to premature termination of the protein. As a consequence, Δ FosB lacks degradation signals present in the C-terminus of FosB. In addition, various phosphorylation events at the N-terminus of Δ FosB also contribute to its stabilization (Nestler, 2013). The stability of Δ FosB has been viewed as an example of the ways in which drug-induced alterations in gene expression can persist for long periods of time after drug withdrawal. Longer term changes in gene transcription can be further propagated by the ability of Δ FosB to induce histone modifications in target

1 Department of Neuroscience, Karolinska Institute, Stockholm, Sweden

2 Peking-Tsinghua Center for Life Sciences, PKU-IDG/McGovern Institute for Brain Research, Peking University School of Life Sciences, Beijing, China

3 Chinese Institute for Brain Research, Beijing, China

*Corresponding author. Tel: +46 5248 7660; E-mail: carlos.ibanez@pku.edu.cn

genes (Teague & Nestler, 2021). Functional studies in transgenic mice using inducible overexpression of Δ FosB or AP1 dominant-negative proteins in medium spiny neurons (MSNs) of the NAc support the notion that accumulation of Δ FosB increases sensitivity to the locomotor and rewarding effects of cocaine and other drugs of abuse, increases cocaine self-administration, and promotes positive reinforcement (Peakman *et al*, 2003; Zachariou *et al*, 2006; Kelz *et al*, 1999). Based on these results, it has been proposed that Δ FosB serves as a critical molecular switch for addiction (Nestler, 2013; Nestler *et al*, 2001; Robison & Nestler, 2022).

Despite the importance of Δ FosB as a key regulator of reward responses, very little is known about the molecular mechanisms that mediate its production by alternative splicing of FosB mRNA. The RNA-binding protein PTB (polypyrimidine tract-binding protein) was found to interact with the 3' end of intron 4 of FosB mRNA, and PTB overexpression in HeLa or PC12 cells decreased the splicing efficiency of a FosB minigene transcript, favoring retention of intron 4 (Marinescu *et al*, 2007; Alibhai *et al*, 2007). These results suggested that PTB may be a negative regulator of Δ FosB production. However, the actual role of this protein in Δ FosB induction is unclear, as PTB was not regulated by either acute or chronic drug administration (Alibhai *et al*, 2007), and splicing of FosB mRNA intron 4 was not affected by PTB knock-down (Marinescu *et al*, 2007). Based on these and other findings, it has been postulated that the generation of Δ FosB mRNA is constitutive and essentially proportional to the levels of FosB mRNA, specially under conditions of high gene induction (Alibhai *et al*, 2007). Accumulation of Δ FosB in NAc during chronic drug exposure is thought to be solely due to the high stability of the protein, not to increased generation of Δ FosB mRNA, and evidence that dopamine signaling can regulate FosB mRNA splicing to contribute to the generation of Δ FosB is currently lacking.

Activin receptor-like kinase 4 (ALK4) is a type I serine–threonine kinase receptor for a subset of TGF β superfamily ligands that include activins A, B, and members of the growth differentiation factor (GDF) subfamily, such as GDF-1, –3 and –5 (Schmierer & Hill, 2007; Moustakas & Heldin, 2009; ten Dijke *et al*, 1994). ALK4 is an essential component of activin A signaling, while activin B, GDF-1, and GDF-3 can signal by either ALK4 or the related receptor ALK7 (Andersson *et al*, 2008, 2006; Tsuchida *et al*, 2004). The main downstream effectors of activated TGF β superfamily receptors are the Smad proteins (Derynck & Budi, 2019). Canonical signaling by ALK4 involves the recruitment and phosphorylation of Smad2 and Smad3, which then partner with Smad4 and translocate to the cell nucleus, where they regulate gene expression in conjunction with cell type-specific transcription factors (Schmierer & Hill, 2007; Budi *et al*, 2017; Massagué, 2012). ALK4 is expressed in the extra-embryonic ectoderm and epiblast of early postimplantation embryos; by midgestation, its expression is nearly ubiquitous, including the central nervous system (Gu *et al*, 1998; Verschuere *et al*, 1995). A null mutation in mouse *Acvr1b*, the gene-encoding ALK4, is embryonic lethal due to defects in primitive streak formation and gastrulation (Gu *et al*, 1998). Using a conditional approach, a recent study from our laboratory demonstrated the role of ALK4 in the regulation of cortical somatostatin interneuron development (Göngrich *et al*, 2020). However, a large gap of knowledge remains on the functions of ALK4 in the nervous system, particularly in the adult brain. Various activities have been attributed to activins and related ligands in nervous tissue, specially in paradigms

of neural injury and morphological plasticity, mainly through gain-of-function studies using exogenous activin administration (Hasegawa *et al*, 2014; Su *et al*, 2018). In the context of drug-induced behaviors, it has been reported that infusion of activin A in the NAc increases the rate of cocaine self-administration in rats (Gancarz *et al*, 2015). However, the role, if any, of endogenous activin/ALK4 signaling in the NAc, or its possible relationship to molecular mechanisms of reward, such as Δ FosB, remains unknown.

In this study, we investigated the contribution of mechanisms operating at the mRNA level to the regulation of Δ FosB production after stimulation of dopamine D1 receptors in primary cultures of NAc MSNs using a novel *in vitro* sensitization paradigm. We discovered a previously unappreciated synergy between dopamine and activin/ALK4 signaling leading to robust generation of Δ FosB mRNA. Further exploration of this finding led to the identification of the RNA-binding protein PCBP1 as the first known positive regulator of FosB mRNA splicing, and validation of its requirement for Δ FosB mRNA generation in response to concurrent dopamine and activin signaling. Strikingly, ablation of ALK4 in NAc MSNs of adult mice completely abolished drug-induced induction of Δ FosB mRNA and behavioral sensitization to cocaine.

Results

Signaling by dopamine D1 and ALK4 receptors synergize to potentiate induction of Δ FosB mRNA and nuclear translocation of Δ FosB protein in NAc MSNs

Sensitization is the phenomenon by which exposure to a stimulus induces a larger response to subsequent exposure. Studies in rats and mice have shown that a single exposure to a psychostimulant, such as cocaine, can induce behavioral sensitization in both a context-dependent and a context-independent manner. Since enhanced responses to the second exposure can be observed days or even weeks after the first, this paradigm has been utilized to study persistent neural plasticity induced by psychoactive drugs and other stimuli. We investigated the effects of stimulation of dopamine receptors in primary cultures of NAc MSNs using an *in vitro* sensitization paradigm consisting of two consecutive treatments with the D1 agonist SKF81297 (Fig 1A). We initially focused on the D1 receptor because of its well-established role in the regulation of Fos family proteins, including FosB (Young *et al*, 1991; Hope *et al*, 1992). A single 2 h stimulation of MSNs with SKF81297 induced a small (1.7-fold) increase in FosB mRNA levels, but no significant change could be observed in c-Fos or Δ FosB mRNAs, nor in the levels of ARC and EGR1 (a.k.a. Ziff268) mRNAs, two other critical mediators of transcriptional responses to cocaine and D1 signaling (Jiang *et al*, 2021; Renthall *et al*, 2009; Penrod *et al*, 2020; Valjent *et al*, 2006) (Appendix Fig S1A–F). Likewise, nuclear translocation of Δ FosB protein was also unaltered by a single stimulation with SKF81297 (Appendix Fig S1G and H). In contrast, two consecutive 2 h treatments separated by 2 days produced a 3-fold increase in Δ FosB mRNA (Fig 1A and B). Induction of FosB mRNA remained at about 2-fold (Fig 1C), while no significant changes were observed in c-Fos, ARC, and EGR1 mRNAs (Fig 1D–F). Stimulation of ALK4 receptors with activin A had no effect on Δ FosB mRNA levels, either after a single treatment or two consecutive exposures (Fig 1B;




Figure 1. Dopamine and ALK4 signaling synergize to potentiate induction of Δ FosB mRNA and nuclear translocation of Δ FosB protein in NAC MSNs.

A Schematic of the protocol used for consecutive stimulation of primary cultures of NAC MSNs. MSNs were extracted at embryonic day 16.5 (E16.5) and cultured for 7 days (7div) before 2 h (2 h) treatment with either 1 μ M SKF81297 (SKF), 100 ng/ml activin A (Act) or their combination (Act+SKF). Following 2 additional days (2div), the treatments were repeated and cultures were fixed for imaging or collected for RNA extraction.

B–F Real-time quantitative PCR (RT-qPCR) analysis of Δ FosB (B), FosB (C), c-Fos (D), ARC (E), and EGR1 (F) mRNAs in MSNs cultures exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Results are expressed as average \pm SEM of fold increase over the vehicle. $N = 5$.

G Representative immunofluorescence staining of Δ FosB (green) in cultured MSNs exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with MAP2 and DAPI is shown in the lower row. Scale bar, 25 μ M.

H Quantification of Δ FosB⁺ nuclei in cultured MSNs after the indicated treatments (same treatment on both treatment days). Results are expressed as average \pm SEM of % of MAP2⁺ cells displaying Δ FosB in the nucleus. $N = 6$.

I Representative immunofluorescence staining of Δ FosB (green) in cultured MSNs exposed to consecutive stimulation, with different 1st and 2nd treatments with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with MAP2 and DAPI is shown in the right column. Scale bar, 25 μ M.

J Quantification of Δ FosB⁺ nuclei in cultured MSNs after different 1st and 2nd treatments with activin A (Act), SKF81297 (SKF), or their combination as indicated. Results are expressed as average \pm SEM of % of MAP2⁺ cells displaying Δ FosB in the nucleus. $N = 6$.

K Representative immunofluorescence staining of Δ FosB (green) in cultured MSNs extracted from mutant mice lacking ALK4 in GABAergic neurons (*Gad67^{Cre}; Alk4^{fl/fl}*) and control mice (*Alk4^{fl/fl}*) after exposure to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with MAP2 and DAPI is shown in the second and fourth rows. Scale bar, 25 μ M.

L Quantification of Δ FosB⁺ nuclei in cultured MSNs from mutant and control mice after the indicated treatments (same treatment on both treatment days). Results are expressed as average \pm SEM of % of MAP2⁺ cells displaying Δ FosB in the nucleus. $N = 6$.

Data information: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; one-way ANOVA followed with the Tukey's multiple comparisons test (B–F, H, J); two-way ANOVA followed with the Tukey's multiple comparisons test (L). N denotes biological replicates.

Appendix Fig S1B). Neither were c-Fos, FosB, ARC, or EGR1 mRNA levels affected by activin A on its own (Fig 1C–F; Appendix Fig S1C and D and G and H). On the other hand, the response of Δ FosB mRNA to two consecutive treatments with SKF81297 was significantly augmented (>10-fold) by co-stimulation with activin A (Fig 1B). Under these conditions, herein termed consecutive co-stimulation, FosB and ARC mRNAs levels increased 3-fold, and EGR1

mRNA 6-fold, while no change was observed in c-Fos mRNA (Fig 1C–F). Importantly, the response of Δ FosB mRNA to consecutive co-stimulation with SKF81297 and activin A was paralleled by induction of Δ FosB protein and its translocation to MSNs nuclei (Fig 1G and H). Although activin A had no effect on its own, it greatly potentiated the effect of SKF81297 on the nuclear translocation of Δ FosB protein (Fig 1H). Neither SKF81297, activin A, or its

combination affected the levels of *Acvr1b* mRNA, encoding ALK4, in MSN primary cultures (Appendix Fig S11).

The changes in responsiveness that accompany sensitization to a second exposure to a psychostimulant are thought to be induced by the first stimulation (“induction phase”), while the second stimulation serves to reveal the underlying sensitization effect (“expression phase”). The strong synergy between SKF81297 and activin A after consecutive co-stimulation prompted us to investigate whether activin A exerted its potentiation effects during the first exposure, the second, or both. In order to test this, activin A was provided along with SKF81297 only during the first (“induction”) or second (“expression”) stimulation of MSN primary cultures. These were compared to consecutive treatments with SKF81297 alone or together with activin A. Only when administered along with SKF81297 in the 1st treatment (induction phase) could activin A potentiate nuclear translocation of Δ FosB protein in MSNs, fully recapitulating the effect of consecutive co-stimulation (Fig 1I and J). No potentiation was observed if activin A was only administered in the 2nd treatment (expression phase), indicating that the potentiation effects of activin A on dopamine receptor signaling take place during the induction phase of sensitization. The effects of two consecutive co-stimulations with SKF81297 and activin A in Δ FosB mRNA induction and Δ FosB protein nuclear translocation were significantly reduced by SB431542, an inhibitor of ALK4, the main activin receptor, and PKI 14–22 amide (PKI14–22), an inhibitor of Protein Kinase A (PKA), a downstream effector of D1 receptor signaling, respectively (Appendix Fig S1J–L).

Acvr1b mRNA, encoding ALK4, was found to be expressed in 70 to 75% of DARPP-32⁺ and D1⁺ neurons in the NAc and striatum (Str) (Appendix Fig S2A–C). These neurons are GABAergic, and *Gad67^{Cre}*; *Alk4^{fl/fl}* mutant mice expressing Cre recombinase in GABAergic neurons showed a significant reduction in *Acvr1b* mRNA levels in the NAc and Str, but less so in other areas, including motor cortex (Appendix Fig S2D–F). ALK4 protein expression, as detected by immunohistochemistry, was also reduced in NAc, but not in the primary motor cortex, of these mutant mice (Appendix Fig S2G and H). Nuclear translocation of Δ FosB protein induced by two consecutive co-stimulations with SKF81297 and activin A was investigated in primary cultures of MSNs isolated from these mutant mice. Activin A was unable to potentiate the effects of SKF81297 in MSNs isolated from *Gad67^{Cre}*; *Alk4^{fl/fl}* mutant mice (Fig 1K and L), although these neurons responded normally to SKF81297 alone (Fig 1L).

Concurrent signaling by D1 and ALK4 receptors induces phosphorylation, nuclear translocation, co-localization, and direct interaction of Smad3 and the RNA-binding protein PCBP1 in NAc MSNs

We considered the possibility that ALK4 signaling may play a role in the generation of Δ FosB mRNA by enhancing the alternative splicing of FosB mRNA. Although Smad proteins were not known to play a role in mRNA splicing, it was shown by Tripathi *et al.*, 2016 that Smad3 could regulate alternative splicing of CD44 mRNA through direct interaction with the RNA-binding protein PCBP1. In addition to Smad3 activation, this process was dependent on concurrent phosphorylation and nuclear translocation of PCBP1 induced by treatment with either epidermal growth factor or serum. The requirement of PKA activity for the synergistic interaction between

D1 and ALK4 receptor signaling in Δ FosB mRNA induction and nuclear translocation of Δ FosB protein (Appendix Fig S1J–L) suggested the possibility that PCBP1 may be a target of PKA-mediated phosphorylation. In line with this notion, two consecutive stimulations (as in Fig 1A) of HeLa cells with forskolin, a strong PKA activator, induced robust phosphorylation of endogenous PCBP1, as detected with antibodies against phospho-PKA substrates, which could be blocked by the PKA inhibitor PKI14-22 (Fig 2A). Moreover, forskolin induced a strong enrichment of PCBP1 in the nuclear fraction of these cells, which could be inhibited by PKI14-22, although it was not significantly affected by activin A or the ALK4 inhibitor SB431542 (Fig 2B). Smad3 and PCBP1 could be co-immunoprecipitated from lysates of HeLa cells after two consecutive co-stimulations with forskolin and activin A but not after treatment with either agent alone (Fig 2C and D). In order to verify these observations in NAc MSNs, we immunostained neuronal cultures for Smad3 and PCBP1 after two consecutive stimulations with SKF81297, activin A or both. As expected, activin A induced robust nuclear translocation of Smad3 in these cultures, while SKF81297 had no effect (Fig 2E and F). On the other hand, SKF81297 induced nuclear translocation of PCBP1 and co-stimulation with activin A produced a small potentiation of this effect (Fig 2E and G). Importantly, two consecutive co-stimulations with SKF81297 and activin A induced a strong co-localization of PCBP1 and Smad3 in MSN nuclei, while individual treatment with either agent had a much lower effect (Fig 2E and H). We note that the level of PCBP1 mRNA in MSNs was not affected by any of these treatments (Fig 2I). To assess direct PCBP1/Smad3 interaction in neurons, we used the Proximity Ligation Assay (PLA) in primary cultures of MSNs treated with SKF81297 and activin A. Two consecutive co-stimulations induced a pronounced increase in PLA signals in MSNs, with little effect of either agent on its own (Fig 2J and K). Treatment with the SB431542 or PKI14-22 inhibitors completely abolished the synergistic effect of SKF81297 and activin A on PCBP1/Smad3 interaction (Fig 2J and K), suggesting that concurrent phosphorylation of PCBP1 and Smad3 by PKA and ALK4 kinases, respectively, is required for their interaction in MSN nuclei.

PCBP1 and Smad3 interact with regulated intron 4 and adjacent sequences in FosB mRNA upon coincident activation of D1 and ALK4 receptors in NAc MSNs

RNA splicing factors may interact directly or indirectly with their target pre-mRNAs but do not typically remain bound to the mature mRNA after splicing. In order to determine whether PCBP1 and Smad3 interact with FosB and Δ FosB mRNAs, we assessed the presence of these transcripts by quantitative RT-PCR in PCBP1 and Smad3 immunoprecipitates from extracts of HeLa cells after two consecutive stimulations with activin A, forskolin or both. No Δ FosB mRNA could be detected under any condition in either PCBP1 or Smad3 immunoprecipitates ($N=3$), suggesting that these proteins do not associate with the Δ FosB transcript. Very low levels of FosB mRNA could be detected in PCBP1 and Smad3 immunoprecipitates of untreated cells, while a moderate increase was observed after either forskolin or activin A treatment, respectively (Fig 3A and B). On the other hand, two consecutive co-stimulations of HeLa cells with forskolin and activin A strongly potentiated the amount of FosB mRNA recovered from both PCBP1 and Smad3




Figure 2.

Figure 2. Concurrent signaling by dopamine and ALK4 induce phosphorylation, nuclear translocation, co-localization, and direct interaction of Smad3 and the RNA-binding protein PCBP1 in NAc MSNs.

- A Western blot analysis of Protein Kinase A phospho-substrates (PKA P -subs) in PCBP1 immunoprecipitates from HeLa cells exposed to sequential stimulation (same treatment on both treatment days) with forskolin (Fsk) in the absence or presence of the PKA inhibitor PKI14-22 (PKI). Reprobing with anti-PCBP1 antibodies is shown in the middle blot. The lower blot shows the total amount of PCBP1 in the cell lysates. Quantification of three independent experiments is shown on the right. Results are expressed as average \pm SEM of fold increase over vehicle normalized to PCBP1 levels.
- B Nuclear localization of PCBP1 analyzed by Western blotting of extracts from HeLa cells exposed to sequential stimulation (same treatment on both treatment days) with activin A (Act), forskolin (Fsk), PKA inhibitor (PKI), and the ALK4 inhibitor SB431542 (SB) as indicated. Blots of nuclear and cytosolic fractions were reprobed with anti-PCNA and anti- α Tubulin (α Tub) antibodies, respectively. Quantification of three independent experiments is shown on the right. Results are expressed as average \pm SEM of fold increase over the vehicle (first column) normalized to PCBP1 levels.
- C Western blot analysis of Smad3 in PCBP1 immunoprecipitates from HeLa cells exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), forskolin (Fsk), or their combination as indicated. The Act+Fsk extracts were also immunoprecipitated using control IgG antibodies as an additional control. Reprobing with anti-PCBP1 antibodies is shown in the middle blot. The lower blot shows unaltered Smad3 levels across all conditions. Quantification of four independent experiments (biological replicates) is shown on the right. Results are expressed as average \pm SEM of fold increase in Smad3/PCBP1 co-immunoprecipitation (coIP) over vehicle (first column) normalized to PCBP1 levels.
- D Western blot analysis of PCBP1 in Smad3 immunoprecipitates from HeLa cells using conditions identical to panel (C). The lower blot shows unaltered PCBP1 levels across all conditions. Quantification of four independent experiments (biological replicates) is shown on the right. Results are expressed as average \pm SEM of fold increase in PCBP1/Smad3 co-immunoprecipitation (coIP) over vehicle (first column) normalized to Smad3 levels.
- E Representative immunofluorescence staining of Smad3 (green) and PCBP1 (red) in cultured MSNs exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with MAP2 and DAPI is shown in the lower row. Scale bar, 25 μ M.
- F, G Quantification of PCBP1⁺ (F) or Smad3⁺ (G) nuclei in cultured MSNs after the indicated treatments (same treatment on both treatment days). Results are expressed as average \pm SEM of % of MAP2⁺ cells displaying PCBP1 (F) or Smad3 (G) in the nucleus. $N = 5$.
- H Quantification of PCBP1/Smad3 co-localization in nuclei of cultured MSNs using conditions identical to panel (F). Results are expressed as average \pm SEM (biological replicates) of % of MAP2⁺ cells displaying co-localization of PCBP1 and Smad3 in the nucleus.
- I RT-qPCR analysis of PCBP1 mRNA in cultured MSNs exposed to consecutive stimulation as indicated (same treatment on both treatment days). Results are expressed as average \pm SEM of fold change over vehicle. No significant differences were found by one-way ANOVA followed by the Tukey's multiple comparisons test ($P \geq 0.05$; $N = 4$).
- J Representative images of PLA (red) between Smad3 and PCBP1 (S3/PCBP1) in cultured MSNs exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF), PKA inhibitor (PKI) or SB431542 (SB) as indicated. Counterstaining with MAP2 and DAPI is shown in the lower row. Scale bar, 25 μ M.
- K Quantification of PLA between Smad3 and PCBP1 in cultured MSNs after the indicated treatments. Results are expressed as average \pm SEM of the number of PLA dots per cell. $N = 5$.

Data information: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; one-way ANOVA followed with the Tukey's multiple comparisons test. N denotes biological replicates. Source data are available online for this figure.

immunoprecipitates (Fig 3A and B), suggesting that the two pathways synergize to induce the association of these proteins with FosB mRNA. This question was addressed further using a modified PLA protocol developed to detect RNA/protein interactions (RNA-PLA) (Roussis *et al*, 2016; Zhang *et al*, 2016). We used antisense probes designed to recognize different regions in FosB mRNA, including a region in exon 4 (pE4) immediately upstream of intron 4 (*i.e.*, the regulated intron), intron 4 (prI4), exon 3 (pE3), and a probe against the exon 4a/4b (pE4a/b) junction that forms after removal of intron 4, which is present in Δ FosB but not in FosB mRNA (Fig 3C). Two consecutive co-stimulations of HeLa cells with forskolin and activin A produced robust interaction of PCBP1 and Smad3 with probes directed to exon 4 and intron 4 of FosB mRNA (Fig 3D–F; Appendix Fig S2A–C) but not with probes directed to exon 3 nor the exon 4a/4b junction (Appendix Fig S3D and E). Stimulation with either forskolin or activin A on their own had little effect (Fig 3D–F; Appendix Fig S3A–C). The interactions of PCBP1 and Smad3 with FosB mRNA sequences were completely eliminated by treatment with PKA or ALK4 kinase inhibitors (Fig 3D–F; Appendix Fig S3A–C). No RNA-PLA signals were detected with sense probes for either exon 4 or intron 4 (Appendix Fig S3D and E), nor with antisense probes and an unrelated antibody (Appendix Fig S3F). Importantly, comparable results for both PCBP1 and Smad3 were obtained in primary cultures of NAc MSNs using the antisense probe against exon 4 (pE4) of FosB mRNA (Fig 3G–I).

PCBP1 is required for the synergistic effects of dopamine and ALK4 signaling on Δ FosB mRNA induction and nuclear translocation of Δ FosB protein in NAc MSNs

In order to assess the requirement of PCBP1 for the induction of Δ FosB mRNA in response to synergistic activation of D1 and ALK4 receptors, we developed a lentiviral vector expressing shRNA targeting PCBP1 mRNA. In primary cultures of MSNs neurons, infection with shPCBP1 lentivirus reduced PCBP1 mRNA expression by 90%, while a lentivirus carrying a scrambled shRNA had no effect (Fig 4A). In agreement with our previous findings, two consecutive co-stimulations of NAc MSNs with SKF81297 and activin A produced a robust induction of Δ FosB mRNA (>17-fold *vs.* vehicle), while SKF81297 on its own had a much lower effect (3-fold) (Fig 4B). Interestingly, knock-down of PCBP1 strongly blunted the response of MSNs to consecutive co-stimulation with SKF81297 and activin A but had no effect on the response to SKF81297 in the absence of activin A (Fig 4B). In contrast, FosB mRNA levels were unaffected by PCBP1 knock-down regardless of treatment (Fig 4C). Knock-down of PCBP1 also eliminated the synergistic effects of SKF81297 and activin A on the nuclear translocation of Δ FosB protein in MSNs (Fig 4D and E), phenocopying the response of neurons lacking ALK4 (Fig 1L). The effect of SKF81297 in the absence of activin A was again unaffected by PCBP1 knock-down (Fig 4E).




Figure 3.

Figure 3. PCBP1 and Smad3 interact with regulated intron 4 and adjacent sequences in FosB mRNA upon coincident activation of D1 and ALK4 receptors in NAC MSNs.

- A, B RT-qPCR analysis of FosB mRNA in PCBP1 (A) and Smad3 (B) immunoprecipitates from HeLa cells exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), forskolin (Fsk) or their combination as indicated. Results are expressed as average \pm SEM of fold increase over vehicle. $N = 4$.
- C Schematic showing the location of probes used for RNA-PLA studies on the interaction of PCBP1 and Smad3 with sequences in FosB and Δ FosB mRNAs. The top diagram shows the FosB gene with the relative position of exons (E1 to E4) and introns (I1 to I4). The location of the regulated intron 4 (rI4) subjected to alternative splicing in FosB mRNA is indicated with black dotted lines. In Δ FosB mRNA, the dotted line indicates the junction between exons 4a and 4b after rI4 splicing. The locations of probes for exon 3 (pE3), exon 4 (pE4), regulated intron 4 (pI4), and exon 4a/4b junction (pE4a/b) are indicated in green. The red bar shows the location of the stop codon that becomes in-frame in Δ FosB mRNA after rI4 splicing.
- D Representative images of RNA-PLA (green) between PCBP1 (left) or Smad3 (right) and sequences in exon 4 (pE4) of FosB mRNA, respectively, in HeLa cells exposed to consecutive stimulation (same treatment on both treatment days) as indicated. Counterstaining with phalloidin and DAPI is shown in the second and fourth columns. Scale bars, 20 μ M.
- E, F Quantification of RNA-PLA between PCBP1 (E) or Smad3 (F) and pE4 in HeLa cells after the indicated treatments. Results are expressed as average \pm SEM of the number of RNA-PLA dots per cell. Only nuclear dots were quantified in this analysis. $N = 4$.
- G Representative images of RNA-PLA (green) between PCBP1 (left) or Smad3 (right) and sequences in exon 4 (pE4) of FosB mRNA, respectively, in cultured MSNs exposed to consecutive stimulation as indicated. Counterstaining with phalloidin and DAPI is shown in the second and fourth columns. Scale bars, 20 μ M.
- H, I Quantification of RNA-PLA between PCBP1 (E) or Smad3 (F) and pE4 in cultured MSNs after the indicated treatments. Results are expressed as average \pm SEM of the number of RNA-PLA dots per cell. Only nuclear dots were quantified in this analysis. $N = 4$.

Data information: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; one-way ANOVA followed with the Tukey's multiple comparisons test. N denotes biological replicates.

ALK4 is required in NAC MSNs of adult mice for behavioral sensitization to cocaine

The above results show that ALK4 and dopamine signaling synergize in an *in vitro* paradigm of sensitization based on primary cultures of NAc MSNs. This effect was manifested by a greatly potentiated induction of Δ FosB mRNA, and nuclear translocation of Δ FosB protein, following two consecutive co-stimulations of ALK4 and D1 receptors. Based on these results, we investigated the importance of ALK4 signaling in an *in vivo* paradigm of sensitization to cocaine known as the “Two-Injection Protocol of Sensitization” or TIPS (Fig 5A). TIPS has been widely used to study behavioral sensitization to a variety of addictive stimuli, including psychoactive drugs, such as amphetamine, morphine, nicotine, and cocaine (Bernardi & Spanagel, 2014; Valjent *et al.*, 2010). Locomotor behavior in response to cocaine injection was first studied in *Gad67^{Cre}*; *Alk4^{fl/fl}* mutant mice. A single injection of cocaine (sensitization) induced a robust locomotor response of similar magnitude in both mutant and control mice compared with animals that received saline, indicating that ALK4 is not required for acute locomotor responses to this drug (Fig 5B, left). Seven days after the first injection, saline and cocaine groups were divided into two subgroups, each of which received a second injection of either saline or cocaine (expression). Consecutive cocaine injections induced a strong potentiation of the locomotor response in control mice, doubling the distance traveled by the mice that received cocaine only the second time (Fig 5B, right), indicative of successful sensitization. In contrast, ALK4 mutant mice showed no potentiation following consecutive cocaine administrations (Fig 5B); *i.e.*, their locomotor response to the second injection was the same regardless whether they received saline or cocaine in the first injection. These data indicate that induction of behavioral sensitization to cocaine requires a functional ALK4 receptor in GABAergic neurons. As a control experiment, targeting ALK4 recombination in GABAergic neurons derived from the medial ganglionic eminence (using *Nkx2.1^{Cre}*; *Alk4^{fl/fl}* mutant mice), which comprise the majority of interneurons in the cortex and hippocampus but not NAc MSNs (Xu *et al.*, 2008), had no effect on *Acvr1b* mRNA expression in the NAc (Appendix Fig S4A), nor did it affect cocaine sensitization and

locomotor responses in the TIPS paradigm (Fig 5C). In order to more specifically assess the importance of ALK4 in NAc MSNs, we used a *Gpr101^{Cre}* transgenic line that specifically targets Cre recombinase to both the D1 and D2 subtypes of these neurons (Reinius *et al.*, 2015). These mutant mice (*Gpr101^{Cre}*; *Alk4^{fl/fl}*) showed a pronounced reduction in *Acvr1b* mRNA expression in NAc and striatum but not in the primary motor cortex (Appendix Fig S4B–D). Similar to *Gad67^{Cre}*; *Alk4^{fl/fl}* mice, *Gpr101^{Cre}*; *Alk4^{fl/fl}* mutant mice lacking ALK4 specifically in NAc MSNs showed no cocaine sensitization; *i.e.*, their locomotor response to a second cocaine injection was the same irrespectively of the nature of the first injection (Fig 5D), strongly arguing for a specific requirement of ALK4 in NAc MSNs for behavioral sensitization to cocaine. In order to evaluate the acute requirement of ALK4 in sensitization to cocaine independently of possible developmental effects, we stereotaxically delivered adeno-associated viruses (AAV2) expressing Cre and GFP, or GFP alone, to the NAc of adult *Alk4^{fl/fl}* mice (Appendix Fig S4E). The CRE-GFP virus reduced the expression of *Acvr1b* mRNA by 60% in the NAc of these mice compared with control virus expressing only GFP (Appendix Fig S4F) but did not affect *Acvr1b* mRNA expression in the motor cortex (Appendix Fig S4G). Importantly, injection of CRE-GFP virus in the NAc of *Alk4^{fl/fl}* mice abolished locomotor potentiation induced by consecutive cocaine administration in the TIPS paradigm, although it had no effect on the acute effects to a single injection (Fig 5E). Control GFP virus did not affect locomotion (Fig 5E). This result demonstrates that ALK4 is required at the time of drug exposure for behavioral sensitization to cocaine.

Activin A is induced in NAc microglia after cocaine TIPS

We assessed the expression of endogenous ALK4 ligands in the NAc and their possible regulation upon cocaine sensitization. We found expression of *Inhba* mRNA (encoding activin A) and *Gdf-1* mRNA (encoding GDF-1) in several brain structures, including NAc/striatum (Appendix Fig 5A and B). In the NAc, activin A immunoreactivity was detected among *DARPP32⁺* MSNs, *GFAP⁺* astrocytes, and *Iba1⁺* microglia cells (Appendix Fig S5C). Strong immunoreactivity for activin A was found in microglia cell bodies of both NAc and striatum (Appendix Fig S5D). A single injection of cocaine produced




Figure 4. PCBP1 is required for the synergistic effects of dopamine and ALK4 signaling on Δ FosB mRNA induction and nuclear translocation of Δ FosB protein in NAc MSNs.

- A RT-qPCR analysis of PCBP1 mRNA in cultures of naive MSNs (no virus) or MSNs treated with lentiviruses expressing shRNA against PCBP1 (shPCBP1) or a scrambled shRNA. Results are expressed as average \pm SEM of fold increase over control uninfected cells, $N = 4$.
- B, C RT-qPCR analysis of Δ FosB (B) and FosB (C) mRNA in cultured MSNs treated with scrambled and shPCBP1 lentiviruses and exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Results are expressed as average \pm SEM of fold increase over vehicle. $N = 4$.
- D Representative immunofluorescence staining of Δ FosB (red) in cultured MSNs treated with scrambled and shPCBP1 lentiviruses and exposed to consecutive stimulation (same treatment on both treatment days) with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with green fluorescent protein (GFP, expressed by the lentiviruses) and DAPI is shown in the second and fourth rows. Scale bar, 25 μ m.
- E Quantification of Δ FosB⁺ nuclei in cultured MSNs treated with scrambled and shPCBP1 lentiviruses followed by the indicated treatments. Results are expressed as average \pm SEM of % of GFP⁺ cells displaying Δ FosB in the nucleus. $N = 4$.

Data information: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; one-way ANOVA followed with the Tukey's multiple comparisons test in (A), two-way ANOVA followed with the Tukey's multiple comparisons test in (B, C, and E). N denotes biological replicates.




Figure 5. ALK4 is required in NAC MSNs of adult mice for behavioral sensitization to cocaine.

A Schematic of the Two-Injection Protocol of Sensitization (TIPS) for the habituation, sensitization, and expression sessions. After entering the activity box, animals remain for 30 min before an intraperitoneal injection of saline. Thirty minutes later, saline (Sal) or cocaine (Coc) are injected after which locomotor activity is recorded for 60 min (red bars). Animals are returned to their home cage at the end of this period. This protocol generates 4 groups of animals, namely Sal/Sal, Sal/Coc, Coc/Sal, and Coc/Coc according to what they received during the 1st and 2nd injections, respectively.

B Locomotor responses of control mice (*Gad67^{Cre}*) and mutant mice lacking ALK4 in GABAergic neurons (*Gad67^{Cre}; Alk4^{fl/fl}*) during cocaine TIPS. “Sensitization” denotes locomotor activity recorded after the first injection of saline (Sal) or cocaine (Coc), while “Expression” denotes locomotor activity in the 4 groups of mice after the second injection. Results are expressed as average ± SEM of distance covered in meters during the 60 min recording period. *N* = 6.

C Locomotor responses of control mice (*Nkx2.1^{Cre}*) and mutant mice lacking ALK4 in GABAergic neurons derived from the medial ganglionic eminence (*Nkx2.1^{Cre}; Alk4^{fl/fl}*) during cocaine TIPS using conditions identical to panel (B). *N* = 6.

D Locomotor responses of control mice (*Gpr101^{Cre}*) and mutant mice lacking ALK4 in NAC MSNs (*Gpr101^{Cre}; Alk4^{fl/fl}*) during cocaine TIPS using conditions identical to panel (B). *N* = 5.

E Locomotor responses of *Alk4^{fl/fl}* mice stereotactically injected in NAC with control AAV2 viruses expressing only GFP (AAV2 GFP) or AAV2 viruses expressing both Cre recombinase and GFP (AAV2 Cre/GFP) during cocaine TIPS using conditions identical to panel (B). *N* = 6.

Data information: Results are expressed as average ± SEM of distance covered in meters during the 60 min recording period. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; two-way ANOVA followed with the Tukey’s multiple comparisons test. *N* denotes biological replicates.

a moderate increase in NAc *Inhba* mRNA, which did not reach statistical significance (Appendix Fig S5E). However, *Inhba* mRNA was significantly upregulated in the NAc of mice that received two consecutive cocaine injections according to the TIPS protocol (Appendix Fig S5F), indicating that cocaine sensitization, but not a single acute exposure, is able to increase *Inhba* mRNA in the NAc. At the protein level, the number of cells expressing activin A increased significantly among the *Iba1*⁺ population (Appendix Fig S5G and H), but not among DARPP32⁺ neurons (Appendix Fig S5I and J) in the mice that received two consecutive cocaine injections. The fact that activin A was specifically regulated by repeated cocaine administration in microglial cells, but not in MSNs or astrocytes, suggests that NAc microglia represent an important source of activin A ligand upon sensitization to cocaine.

ALK4 is required in NAc MSNs of adult mice for sustained induction of Δ FosB mRNA during cocaine sensitization

As expected, acute induction of immediate early genes FosB and c-Fos could be readily detected in NAc of both control (*Gad67*^{Cre}) and *Gad67*^{Cre}; *Alk4*^{fl/fl} mutant mice 1 h after the second cocaine injection (Fig 6A and B). Similar results were observed in *Gpr101*^{Cre}; *Alk4*^{fl/fl} mice (Appendix Fig S6A and B). However, the levels of Δ FosB mRNA remained unchanged at this time point regardless of treatment or genotype (Fig 6C; Appendix Fig S6C). Expression of FosB and c-Fos mRNAs returned to basal levels 24 h after the second injection in the NAc of both control and mutant mice (Fig 6D and E and Appendix Fig S6D and E). In contrast, a strong induction of Δ FosB mRNA was observed 24 h after the second injection in the NAc of control mice that received two consecutive cocaine doses (Fig 6F; Appendix Fig S6F). Importantly, this response was completely abolished in both lines of mutant mice (Fig 6F; Appendix Fig S6F). Similar results were observed at the protein level. Immunoreactivity for Δ FosB was greatly elevated in the NAc of control mice 24 h after two consecutive cocaine injections, but no increase was observed *Gad67*^{Cre}; *Alk4*^{fl/fl} mutant mice (Fig 6G and H). Analysis of NAc lysates by Western blotting confirmed these results, showing a pronounced increase in Δ FosB protein levels in control but not mutant mice 24 h after two consecutive cocaine doses (Fig 6I and J). At this time, the levels of NAc FosB protein remained unchanged in both strains (Fig 6K). Similar to Δ FosB, the levels of ARC and EGR1 mRNAs were also greatly induced 24 h after the second cocaine injection (Appendix Fig S7A–D). However, mutant mice showed no lasting induction of ARC and EGR1 mRNAs 24 h after cocaine TIPS (Appendix Fig S7A–D). Together, these results indicate a profound deficit in long-lasting induction of Δ FosB and its downstream targets ARC and EGR1 in NAc after specific ablation of ALK4 in MSNs.

Discussion

Since its discovery three decades ago (Mumberg *et al*, 1991; Naka-beppu & Nathans, 1991), Δ FosB has been the subject of numerous investigations, with hundreds of papers describing various aspects of its induction, expression, and downstream effects. The relationship between Fos family proteins and activation of D1 receptors is equally old (Young *et al*, 1991; Hope *et al*, 1992); and the

association of Δ FosB with chronic cocaine administration was also uncovered at about the same time (Hope *et al*, 1994). Despite this large body of research, the molecular mechanisms that promote the alternative splicing of FosB mRNA to generate Δ FosB have been unknown. The current view is that Δ FosB mRNA is generated constitutively above a certain level of FosB mRNA expression (Alibhai *et al*, 2007). According to this model, the machinery that promotes the alternative splicing of FosB mRNA is not thought to be affected or regulated by dopamine signaling. In the present study, we describe a previously unknown mechanism by which dopamine signaling synergizes with the TGF β superfamily receptor ALK4 to potentiate the production of Δ FosB in MSNs of the NAc through activation of the RNA-binding protein PCBP1. In the absence of ALK4, locomotor responses to the second dose of cocaine were identical to the first, and long-lasting Δ FosB mRNA and protein expression were abolished, indicating that persistent expression of Δ FosB mRNA and behavioral sensitization to cocaine are both dependent on ALK4 signaling in NAc MSNs of adult mice. We note that the approaches used in our study to inactivate ALK4 expression in the mouse NAc did not distinguish between the D1 and D2 subtypes of MSNs, which have been shown to mediate distinct responses to repeated cocaine intake (Kravitz *et al*, 2012; Bock *et al*, 2013). Although our studies in MSN cultures using SKF81297 are in line with the involvement of the D1 subtype, the role of ALK4 signaling in D2-expressing MSNs and its possible impact on cocaine sensitization remains to be investigated. Given the critical role of Δ FosB in reward responses and addiction, our findings establish a new paradigm for understanding the early molecular events leading to the structural and functional adaptations that underlie those processes.

The molecular mechanisms that mediate the alternative splicing of FosB mRNA to generate Δ FosB are understudied and remain elusive. PCBP1 (poly-(rC)-binding protein 1, also known as hnRNPE1) has been known to participate in various mRNA processing steps, including alternative splicing, stability, and translational repression (Meng *et al*, 2007; Hwang *et al*, 2017). Phosphorylation of PCBP1 at Thr⁶⁰ and Thr¹²⁷ by Pak1 was shown to induce its translocation to the nucleus, where it regulated the alternative splicing of CD44 mRNA, promoting the inclusion of its variable exon (Meng *et al*, 2007). Subsequent studies found that PCBP1 can be phosphorylated at these and other sites by multiple Ser-Thr kinases, including AKT, CDKs, and MAPKs, following activation of growth factor receptors (Chaudhury *et al*, 2010; Tripathi *et al*, 2016). In conjunction with TGF β signaling, activated PCBP1 and Smad3 were shown to form a complex that translocated to the cell nucleus where it promoted exclusion of CD44 mRNA regulated exon, contributing to tumor formation (Tripathi *et al*, 2016). In the present studies, we found that PCBP1 and Smad3 can be activated through phosphorylation by PKA and ALK4, respectively, inducing their association and nuclear translocation in NAc MSNs, thereby extending this paradigm to D1 and ALK4 receptor signaling in neuronal cells (Appendix Fig S8). The ability of activin A to potentiate PCBP1 nuclear translocation induced by D1 receptor activation, suggests that activated Smad3 and PCBP1 first interact in the cytoplasm, from where Smad3 nuclear localization signals may then facilitate the nuclear translocation of the complex.

We found that PCBP1 and Smad3 interacted with FosB mRNA at and around intron 4 sequences, but not other regions in FosB mRNA, nor the exon 4a/4b junction present in Δ FosB mRNA but absent in




Figure 6.

Figure 6. ALK4 is required in NAc MSNs of adult mice for sustained induction of Δ FosB mRNA and protein during cocaine sensitization.

- A–F RT-qPCR analysis of FosB (A, D), c-Fos (B, E), and Δ FosB (C, F) mRNAs in NAc of *Gad67^{Cre}*; *Alk4^{fl/fl}* (A–C) mutant mice 1 h (A–C) or 24 h (D–F) after the second saline (Sal) or cocaine (Coc) injection of TIPS. Results are expressed as average \pm SEM of fold increase over the saline group (Sal/Sal) of *Gad67^{Cre}* mice. $N = 5$.
- G Representative immunofluorescence staining of Δ FosB in NAc of control (*Gad67^{Cre}*) and mutant (*Gad67^{Cre}*; *Alk4^{fl/fl}*) mice, 24 h after the second injection of the TIPS protocol. Scale bars 20 μ m.
- H Quantification of Δ FosB positive cells 24 h after the second injection of TIPS protocol in the NAc of *Gad67^{Cre}* and *Gad67^{Cre}*; *Alk4^{fl/fl}* mice. Results are expressed as average \pm SEM of the number of Δ FosB⁺ cells per mm². At least 10 images from both left and right hemispheres were analyzed per mouse and condition. $N = 5$.
- I Western blot analysis of FosB and Δ FosB in NAc of *Gad67^{Cre}* and *Gad67^{Cre}*; *Alk4^{fl/fl}* mice 24 h after the second injection of TIPS. Blots were probed with an antibody that recognizes both FosB (upper band, 48 kDa) and Δ FosB (lower band, 38 kDa). The lower blot shows reprobing with anti- β -actin.
- J, K Quantification of FosB (J) and Δ FosB (K) levels in NAc of *Gad67^{Cre}* and *Gad67^{Cre}*; *Alk4^{fl/fl}* mice 24 h after the second injection of TIPS. Results are expressed as average \pm SEM of fold increase in FosB and Δ FosB over the saline group (Sal/Sal) of *Gad67^{Cre}* mice normalized to β -actin levels. $N = 4$ s.

Data information: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; one-way ANOVA followed with the Tukey's multiple comparisons test. N denotes biological replicates. Source data are available online for this figure.

FosB, suggesting the involvement of these proteins in FosB intron 4 splicing rather than other processes, such as Δ FosB mRNA stabilization. We note that, although activin A greatly potentiated the effects of SKF81297 on Δ FosB mRNA induction and Δ FosB nuclear translocation, SKF81297 was also able to enhance both these processes on its own, albeit at more moderate levels. In contrast to the co-stimulation regime, however, SKF81297 induced FosB and Δ FosB expression to comparable degrees when administered in the absence of activin A, in agreement with the prevalent model of constitutive Δ FosB generation proportionally to FosB mRNA (Alibhai et al, 2007; Nestler, 2008, 2013). On the other hand, the fact that PCBP1 knock-down in MSNs specifically interfered with the potentiated induction of Δ FosB mRNA afforded by co-stimulation with activin A and SKF81297 without affecting FosB mRNA levels, suggests a direct role for PCBP1 in enhancing Δ FosB mRNA generation independently of FosB gene transcription. In agreement with this, PCBP1 knock-down did not impact the effects of SKF81297 on Δ FosB expression and nuclear localization in the absence of activin A, indicating that PCBP1 specifically participates in the synergy between D1 with ALK4 signaling that enhances Δ FosB mRNA levels, rather than the basal mechanism of Δ FosB mRNA generation. Similar to its role in the alternative splicing of CD44 mRNA (Meng et al, 2007; Tripathi et al, 2016), the PCBP1/Smad3 complex may regulate the function of components of the basal splicing machinery, such as factors associated with the U2 small nuclear ribonucleoprotein complex, to enhance splicing of FosB mRNA intron 4, thereby increasing the generation of Δ FosB mRNA.

Psychostimulants of abuse, including cocaine, methamphetamine, and ecstasy, have been shown to induce neuroinflammation in the brain through activation of resident endothelial, glial and immune cells (Clark et al, 2013; Ersche & Döflinger, 2017). *In vitro* and *in vivo* studies have provided evidence indicating that repeated exposure to cocaine can induce activation of microglial cells, particularly in striatum, through a mechanism involving increased production of reactive oxygen species (ROS) and inflammasome activity (Burkovetskaya et al, 2020; Chivero et al, 2021; Liao et al, 2016). However, whether microglia activation in turn contributes to reinforce cocaine sensitization is unknown. In our studies, we found significantly increased activin A production in NAc microglia following two consecutive cocaine exposures but not after a single administration. An earlier study reported elevated levels of activin A in the NAc of rats after a cocaine relapse (Wang et al, 2017). Increased activin A production in NAc microglia after repeated cocaine exposure suggests its upregulation may be a consequence of cocaine-

mediated neuroinflammation and suggests a previously unappreciated role for these cells in cocaine sensitization.

In conclusion, we propose that the synergistic effects of D1 and ALK4 signaling mediated by concurrent activation of PCBP1 and Smad3 in NAc MSNs represent a previously unrecognized coincidence-detection mechanism that potentiates Δ FosB production and reward-related behavior in response to repeated exposure to particularly salient stimuli. We postulate that such a mechanism facilitates sensitization to specific stimuli only when these are of sufficient magnitude, duration, and persistence to activate concurrent D1 and ALK4 signaling in NAc MSNs. Since Smad3 and PCBP1 are widely expressed in the brain, they may also be involved in Δ FosB expression induced by other stimuli in other brain regions. Targeting the key components of this mechanism may lead to interventions that counteract the effects of drugs of abuse.

Materials and Methods

Mice

Mice were housed 2–5 per cage in standard conditions in a temperature and humidity-controlled environment, on a 12/12 h light/dark cycle with *ad libitum* access to food and water. The following transgenic mouse lines were used for experiments: *Gad67^{Cre}* (Tolu et al, 2010), *Nkx2.1^{Cre}* (Xu et al, 2008), *Gpr101^{Cre}* (Reinius et al, 2015), and *Alk4^{fl/fl}* (Göngrich et al, 2020). All animals were bred in a C57BL/6J background (Jackson). Both males and females were used in the studies; mice were 12 to 14 weeks old at the start of the experiments. Animal protocols were approved by Stockholms Norra Djurförsöksetiska Nämnd and are in accordance with the ethical guidelines of Karolinska Institutet (ethical permits N173-15; 11,563–2018 and 11,571–2018 and N68/16 following EU directive 2010/63/EU).

Cell culture and treatments

Medium Spiny Neuron (MSN) culture was performed according to Penrod et al, 2011 (Penrod et al, 2011) with minor modifications. Briefly, pregnant dams were killed by cervical dislocation at 16.5 days of gestation, embryos were collected and brains were removed in ice-cooled Ca²⁺/Mg²⁺-free Hank's balanced salt solution (Gibco) with 10 mM HEPES (ThermoFisher). After removal of all meninges, ganglionic eminences, i.e., protostriatum, were dissected in fresh Hank's/HEPES solution. Collected tissue was transferred to 15 ml tubes and

digested with 0.25% Trypsin–EDTA (Gibco) for 30 min at 37°C. Trypsin was inactivated with 25% Newborn Calf Serum (NCS, Gibco) and cells were dissociated by 10 passages through a glass pipette in the presence of 25 U DNase (Roche). Cells were then collected by centrifugation for 5 min at 500xg, resuspended in a plating medium: MEM containing Earl's salts (Gibco), supplemented with 10 mM HEPES, 10 mM sodium pyruvate, 0.5 mM glutamine, 12.5 μ M glutamate, 0.6% glucose and 10% NCS. Cells were centrifuged one more time and again resuspended in the plating medium. Cells were then plated at a density of 30,000 cells per coverslip in 24-well plates; coverslips were coated with 100 μ g/ml poly-D-lysine (Sigma-Aldrich) and 4 μ g/ml mouse Laminin (Cultrex, RnD). One to 3 h later, after cells had adhered to the coverslips, the plating medium was replaced by a conditioned growth medium: Neurobasal (Gibco), containing B27 supplement (Gibco) and 0.5 mM glutamine (Gibco), which had been conditioned on a monolayer of glial cells (see below). One-half of the medium was replaced after 3 days with fresh conditioned growth media. Cells were maintained for 7 days at 37°C in 95% O₂/5% CO₂ atmosphere before the start of the experiments.

For the preparation of glial cell-conditioned medium, glia cultures were prepared from cortices of postnatal day 1–4 (P1-4) mice with the striatal tissue removed. Minced tissue was incubated in Hank's balanced salt solution supplemented with 0.25% Trypsin–EDTA and DNase at 37°C for 30 min. After digestion, an equal volume of glia plating medium (MEM with 10 mM HEPES, 1 mM sodium pyruvate, 0.2 mM glutamine, 10% NCS, 0.6% glucose, and 1X penicillin–streptomycin) was added to inhibit the trypsin and the tissue was collected by centrifugation at 500xg for 5 min. The tissue was resuspended in a glia plating medium, mechanically dissociated using a flame-polished glass pipette, and filtered through a 0.7 μ m cell strainer (Fisher scientific). Cells from one pup were plated onto 10 cm standard tissue culture dishes. Media was replaced one day after plating and once per week for each subsequent week. The conditioned medium was harvested after cell monolayers reached 70% confluence.

HeLa cells were grown in MEM supplemented with 2 mM glutamine (Gibco), 20 μ g/ml penicillin/streptomycin (Gibco), and 10% heat-inactivated fetal bovine serum (FBS, Gibco). Cells were maintained at 37°C in a 95% O₂/5% CO₂ atmosphere. Cell line experiments were performed after cell monolayers reached 80% confluence.

MSN cultures (after 7 days *in vitro*) or cell lines (after 80% confluence) were treated for 2 h with one or more of the substances, as indicated in the Results section, at the following concentrations: Activin A (RnD) 100 ng/ml, SKF91297 (Tocris) 1 μ M, Forskolin (Tocris) 25 μ M, SB41542 (Sigma-Aldrich) 10 μ M, PKI14,22 amide (Tocris) 1 μ M. For the sequential stimulation protocol, cells were washed after 2 h treatment in phosphate buffer saline (PBS, Gibco) and fresh growth media was added. Two days later, cells were treated again for 2 h and washed before fixation or lysis.

RNA extraction and quantitative real-time quantitative PCR (RT-qPCR)

Total mRNA was isolated from cell monolayers and dissected brain areas using RNeasy Mini Kit (Qiagen) according to the manufacturer's protocol. QIASHredder (Qiagen) was used for tissue homogenization prior to RNA extraction. RNA purity and quantity of RNA were measured in a Nanodrop 1000 (ThermoScientific). cDNA was synthesized by reverse transcription using 150 ng of RNA in a

reaction volume of 20 μ l using High Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to the manufacturer's protocol. cDNA samples were diluted 1:3 in distilled water before use. For measurement of 18S RNA levels, a further 1:800 dilution was made. The geometric mean of mRNAs encoding the ribosomal protein 18S was used for normalization. RT-qPCR was performed on the 7500 Real-Time PCR System (Applied Biosystems) with SYBR Green fluorescent probes and the following conditions: 40 cycles of 95°C for 15 s, 60°C for 1 min, and 72°C for 30s. Forward and reverse primers were used at 100 nM each. The primer pairs used were as follows: m18s (Fw: 5'-GCAATTATCCCCATG AACG-3'; Rv: 5'-GGCCTCACTAAAACCATCCAA-3'); h18s (Fw: 5'-CTCAACACGGGAAACCTCAC-3'; Rv: 5'-CGCTCCACCAACTAAGAA CG-3'); Δ FosB (Fw: 5'-AGGCAGAGCTGGAGTCGGAGAT-3'; Rv: 5'-GCCGAGGACTTGAACCTCACTCG-3'); FosB (Fw: 5'-GTGAGAGAT TTGCCAGGGTC-3'; Rv: 5'-AGAGAGAAGCCGTCAGGTTG-3'); c-Fos (Fw: 5'-GGAATTAACCTGGTCTGGA-3'; Rv: 5'-AGAGAGAAGCC GTCAGGTTG-3'); ARC (Fw: 5'-CTTGCCTCTGCTCTGAGC-3'; Rv: 5'-AGAAAGCAGCAGCAAGATGG-3'); EGR1 (Fw: 5'-AATCCCCCTTC GTGACTACC-3'; Rv: 5'-GCAAGATGGAGGGTTGG-3'); Acvr1b (Fw: 5'-CCAAGTGGTGGCAGAGTAT-3'; Rv: 5'-CTGGGACAGAGTCTTCT TGATG-3'); PCBP1 (Fw: 5'-ATAGATGATTGGAACCAGGTAAGT-3'; Rv: 5'-GCTCCAAGGGAGCAAGCGAGCAGT-3'); Inhba (Fw: 5'-AT CATCACCTTTGCCGAGTC-3'; Rv: 5'-ACAGGTCACTGCCTTCCTTG-3'); Gdf-1 (Fw: 5'-TTCTGCCAGGGCACGTGCG-3'; Rv: 5'-GGAGCA GCTGCGTGCATGAG-3'). Each experiment was carried out on biological triplicates for each data point. Data analysis was performed using the 2^{- Δ Ct} method, and relative expression levels were calculated for each sample after normalization against 18S RNA levels.

Co-Immunoprecipitation, nuclear fractionation, Western blotting, and RNA immunoprecipitation

HeLa cell lysates were prepared on ice using RIPA buffer (ThermoFisher) containing protease and phosphatase inhibitors (Roche). Briefly, growth media was removed, cells were washed twice with PBS, then lifted in RIPA buffer using a cell scraper. The samples were centrifuged for 10 min at 5,000 g and the supernatant was collected. Immunoprecipitation from cell lysates was performed using Dynabeads Protein A (ThermoFisher) following the manufacturer's protocol. Protein concentration was measured with the Pierce™ BCA Protein Assay Kit (ThermoFisher), and 200 μ g of total protein lysate was used for each immunoprecipitation reaction. Rabbit anti PCBP1 (1:150, #85341, Cell signaling) and rabbit anti Smad3 (1:100, #9624, Cell signaling) were used for immunoprecipitation. Rabbit anti-c-Jun (1:150, #9165, Cell Signaling) was used as a mock control antibody. For Western blotting, immunoprecipitates were boiled at 95°C for 10 min in sample buffer (Life Technologies) before separation by SDS–PAGE in 4–15% Mini-PROTEAN® TGX™ Precast Gel (Bio-Rad). Proteins were transferred to polyvinylidene fluoride (PVDF) membranes (Amersham). Membranes were blocked with 5% nonfat milk in TBST and incubated with antibodies overnight in 1% milk. The antibodies used were as follows: mouse anti PCBP1 (1:400, sc-137,249 Santa Cruz biotech), rabbit anti Smad3 (1:1,000, #9513, Cell Signaling) and rabbits anti-phospho-PKA substrates (RRXS*/T*) (1:1,000, #9624; Cell Signaling).

Cytosolic and nuclear protein fractions were prepared using NE-PER™ Nuclear and Cytoplasmic Extraction Reagents (ThermoFisher)

following the manufacturer's protocol. Western blotting was performed as indicated above using 20 µg of protein per sample. The antibodies used on membranes containing cytosolic and nuclear protein fractions were: rabbit anti PCBP1 (1:1,000, #8534, Cell Signaling), mouse anti PCNA (1:1,000, MAB424R, Sigma-Aldrich), α Tubulin (1:2,000, 05-829, Sigma-Aldrich). For FosB/DFosB analysis in brain tissue mouse-anti FosB (1:1000, Ab11959, Abcam), and Rabbit Anti-b-Actin (1:1000, 4967, Cell Signaling) were used. Immunoreactivity was detected using appropriate horseradish peroxidase (HRP)-conjugated secondary antibodies (1:5,000, Dako). Immunoblots were developed using the Clarity Western ECL substrates (Bio-Rad) or the SuperSignal West Femto Maximum sensitivity substrate (ThermoFisher) and images were acquired with ImageQuant LAS4000 (GE Healthcare). Image analysis and quantification of band intensities were done with ImageQuant software (GE Healthcare).

RNA immunoprecipitation was performed using the Magna RIP® RNA-Binding Protein Immunoprecipitation Kit (Sigma-Aldrich) according to the manufacturer's protocol. Approximately 2×10^7 HeLa cells were used for each immunoprecipitation. Mouse anti PCBP1 (5 µg, sc-137,249 Santa Cruz Biotech) and anti-Smad3 (1:100, #9513, Cell Signaling) were used for immunoprecipitation as described above. Binding to Δ FosB or osSB mRNA was assessed by RT-qPCR as described above using the following primers: Δ FosB (Fw: 5'-AGGCAGAGCTGGAGTCGGAGAT-3'; Rv: 5'-GCCGAGGACTTGAACCTCACTCG-3'); FosB (Fw: 5'-GTGAGAGATTTGCCAGGGTC-3'; Rv: 5'-AGAGAGAAGCCGTCAGGTTG-3').

Immunocytochemistry and proximity ligation assay (PLA)

Coverslips containing cell monolayers were washed twice in PBS for 10 min and fixed for 15 min in 4% PFA (Histolab) / 4% Sucrose. Cells were washed 2 extra times to remove any remaining PFA. Cells were permeabilized in 0.3% Triton X-100 (Sigma-Aldrich) in PBS for 10 min, followed for 1 h blocking in 5% Normal Donkey Serum (NDS, Jackson ImmunoResearch) and 0.05% Triton X-100 in PBS. Cells were then incubated overnight at 4°C under gentle agitation with the primary antibodies in 2% NDS and 0.05% Triton X-100 in PBS. Primary antibodies used were as follows: rabbit anti Δ FosB (1:500, #14695, Cell Signaling), mouse Anti-PCBP1 (1:400, sc-137249 Santa Cruz biotech), rabbit anti Smad3 (1:250, #9513, Cell Signaling), chicken anti MAP2 (1:5,000, ab5392, Abcam) or mouse anti MAP2 (1:200; M4403; Sigma-Aldrich). Coverslips were subsequently washed 3 times for 10 min and incubated with the appropriate secondary antibody for 2 h at room temperature. Secondary antibodies were donkey anti-rabbit IgG Alexa Fluor 555, A31572; donkey anti-goat IgG Alexa Fluor 488, A11055; donkey anti-mouse IgG Alexa Fluor 488, A21202; donkey anti-mouse IgG Alexa Fluor 555, A31570; donkey anti-mouse IgG Alexa Fluor 647, A31571; donkey anti-goat IgG Alexa Fluor 555, A21432 (1:800, Life Technologies and Invitrogen); and donkey anti-chicken Alexa Fluor 647 (1:750; 703-606-155, Jackson ImmunoResearch).

PLA was performed according to the manufacturer's protocol (Sigma-Aldrich). Briefly, cell cultures were fixed as indicated above. Cells were washed 10 min in 0.2% Triton X-100 and blocked for 1 h in 5% NDS and 0.05% Triton X-100 in PBS before incubation with rabbit anti Smad3 (1:250, #9624, Cell Signaling), mouse anti PCBP1 (1:200, sc-137,249; Santa Cruz biotech) and chicken anti MAP2 (Abcam, ab5392, 1:2,000) overnight at 4°C with gentle agitation. Coverslips were subsequently washed with PBS, incubated with minus (anti-mouse)

and plus (anti-rabbit) PLA probes for 1 h at 37°C, washed with buffer A, incubated with Ligation-Ligase solution for 30 min at 37°C, washed again with buffer A, and finally incubated with Amplification-Polymerase solution (Duolink™ *In Situ* detection reagents Orange) for 100 min at 37°C. Alexa anti-chicken was included during the amplification step to reveal the MAP2 signal. After washing with buffer B, sections were incubated for 10 min with DAPI (1:10,000 in buffer B) and mounted on glass slides with DAKO mounting media.

Our RNA-PLA protocol was adapted from Roussis *et al.* (2016) and Zhang *et al.* (2016). One of the primary antibody/probe pairs of standard PLA was replaced by a DNA oligonucleotide containing a 40–50 nucleotide (nt) segment complementary to the RNA of interest, 20–35 nt poly-A as a linker segment, and a common 25 nt segment that serves as one of the PLA hybridization/ligation arms (PLA PLUS probe). Briefly, cells were washed twice for 10 min in Dulbecco's modified Phosphate Saline Buffer (DPBS, ThermoFisher) and fixed with 4% PFA/4% sucrose (Sigma-Aldrich) for 15 min. After two washes in DPBS, cells were permeabilized for 10 min with 0.2% Triton-X100, washed once in DPBS, and blocked in Blocking Buffer (10 mM Magnesium-Acetate, 50 mM Potassium acetate, 250 mM NaCl, 2% BSA, 0.05% Tween-20, 20 µg/ml Sheared Salmon Sperm DNA (ThermoFisher) for 1 h. Probes (100 nM) were heated to 70°C for 5 min in blocking buffer and incubated with the samples overnight at 58°C with gentle agitation. The following probes were used: Exon 4: Antisense: 5'-GGCCCTCTTCGTAGGGGATCTTGCAGCCCGTTTGTGGGCCACC-3'; Sense: 5'-GGTGGCCACAAACCGGCTGCAAGATCCCCTACGAAGAGGGGCC-3'. Regulated Intron 4: Antisense: 5'-CCGAAGCCGCTTCTTACGGGATGTTGACCCTGGCA-3'; Sense: 5'-TGCCAGGGTCAACATCCGCTAAGGAAGACGGCTTCGG-3'. Exon 3: Antisense 5'-TCTCTGCGAACCTTCGCTTTTCTTCTTCTGCGGTA-3'. Exon 4a/b junction: Antisense: 5'-GGAAGGGTCCGCGAGGACTTGAACCTCGGCCAGCGGGCCTGGCCC-3'. All sequences were followed at the 3' with the PLA-chimera probe sequence: 5'-AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATATGACAGAACTAGACTCTT-3'. After overnight incubation, cells were washed three times in DPBS and incubated in NDS containing 0.02% Triton-X100 for 1 h. Primary antibodies were added in 2% NDS, 0.02% Triton-X100, and incubation proceeded at 4°C overnight with gentle agitation. Primary antibodies were used for RNA-PLA were: mouse Anti-PCBP1 (1:200, sc-137,249 Santa Cruz biotech), rabbit anti Smad3 (1:250, #9513, Cell Signaling) and Sirt1 (1:250, #2028; Cell Signaling). Chicken anti MAP2 (1:5,000, ab5392, Abcam) or mouse anti MAP2 (1:200; M4403; Sigma-Aldrich) were used for counterstaining of MSNs; Alexa Fluor A488 Phalloidin (1:1,000, A12379, ThermoFisher) or Alexa Fluor 568 (1:1,000, A12380, ThermoFisher) were used for counterstaining of HeLa cells. After overnight incubation, samples were washed twice with DPBS for 10 min. PLA reactions were then performed as described above.

PCBP1 knock-down by lentivirus infection

Short interference mouse PCBP1 RNAi plasmids were purchased from Origene (TL502540). Lentiviral particles expressing the PCBP1 shRNAi or scramble control were produced by the VirusTech Core Facility, Department of Medical Biochemistry and Biophysics (MBB) Karolinska Institutet. Neurons were infected after 7 days *in vitro* with PCBP1-shRNAi or scrambled sc-shRNAi (MOI: 100) overnight, supplemented with 1 µg/ml polybrene (Sigma-Aldrich).

Two-Injection Protocol of Sensitization (TIPS) to cocaine

TIPS was performed according to Valjent *et al* (2010) (Valjent *et al*, 2010) with minor modifications. Briefly, mice were placed in the center of a 48 d7 48 cm transparent acrylic box equipped with light-beam strips (TSE Actimot system) and allowed to move freely. Distance traveled was calculated using Actimot Software. All mice were habituated to the test apparatus, handling, and procedure for 3 consecutive days before the actual experiment. During the habituation procedure mice were placed for 30 min in the activity box, received a first injection of saline (0.9% NaCl; Sal), were placed back in the box for 30 min, received a second saline injection, and were placed again for 1 h in the box before returning to the home cage. For the first drug injection (day 4, “sensitization”), a similar handling was repeated, except that, the second saline injection was replaced by either a saline or 20 mg/kg cocaine-HCl (Coc; Sigma-Aldrich), resulting in 2 different groups of mice, namely Sal and Coc, respectively. The mice were then placed back in the activity box for 1 h to register their locomotor activity. Seven days later, Sal and Coc groups were further divided before receiving a second injection of saline or cocaine (day 11, “expression”), resulting in 4 groups of mice, namely Sal/Sal, Sal/Coc, Coc/Sal, and Coc/Coc, respectively. The mice were then placed back in the activity box for 1 h to register their locomotor activity and sacrificed after either 1 or 24 h for immunohistochemistry or RNA analysis, as indicated.

Stereotaxic injection of AAV2 vectors

Mice were anesthetized with isoflurane (1–5%) and placed in a stereotaxic frame (David Kopf Instruments). Injection was performed with a Wiretrol capillary micropipette (Drummond scientific) at a flow rate of 50 nl/min (Micro4 controller, World Precision Instruments; Nanojector II, Drummond Scientific). The pipette was left in place for 10 min after the injection before retracting slowly from the brain. *Alk4^{fl/fl}* mice were injected bilaterally with 300 nL of AAV2 adeno-associated viruses expressing either Cre recombinase and GFP (pAAV.CMV.HI.eGFP-Cre.WPRE.SV40) or only GFP (pAAV.CMV.PI.eGFP.WPRE.bGH) (Penn Vector Core, 2.27×10^{13} genomic copies per mL) using coordinates obtained from Paxinos and Franklin atlas as follows: AP +1.7 mm, mediolateral ML ± 0.6 mm, dorsoventral DV -4.5 mm). To evaluate the infection efficiency of the vectors, RNA was extracted and *Acvr1b* mRNA (encoding ALK4) was measured by RT-qPCR (Appendix Fig S4F and G).

Immunohistochemistry, *in situ* hybridization, and image analysis

Mice were deeply anesthetized with isoflurane (Baxter Medical AB) and perfused transcardially with 25 ml of 0.125 M PBS (pH 7.4, Gibco) and 40 ml of 4% paraformaldehyde (PFA, Histolab Products AB). Brains were removed, postfixed in PFA overnight, and cryoprotected in 30% sucrose in PBS. 30 μ m brain sections were cut in a microtome (Leica SM2000 R). Sections were stored at -20 °C in a cryoprotective solution containing 1% DMSO (Sigma-Aldrich) and 20% glycerol (Sigma-Aldrich) in 0.05 M Tris-HCl pH 7.4 (Sigma-Aldrich). For RNA *in situ* hybridization, washing and fixation solutions were prepared using RNAase-free components; surgical

instruments and microtome blades were cleaned with RNaseZap (Sigma-Aldrich) before use.

For immunohistochemistry, brain sections were washed in PBS and blocked for 1 h in 5% normal donkey serum (NDS, Jackson ImmunoResearch) and 0.3% Triton X-100 (Sigma-Aldrich) in PBS. Incubation with primary antibodies was done overnight at 4 °C in a blocking solution. After three consecutive 10 min washes with PBS, sections were incubated with the corresponding secondary donkey Alexa fluorescent antibodies (1:1,000, ThermoFisher) and 0.1 mg/ml of 40-6-diamidino-2-phenylindole (DAPI; Sigma-Aldrich) for 2 h at room temperature. Sections were finally washed with PBS (3×10 min), mounted on glass slides in 0.2% solution of gelatin (Sigma-Aldrich) in 0.05 M Tris-HCl buffer (pH 7.4), dried, and mounted with glass coverslips using DAKO fluorescent mounting medium. The following primary antibodies were used: rabbit anti ALK4 (1:500; ab109300, Abcam); mouse anti DAPP32 (1:750, sc-271,111, Santa Cruz Biotech); rabbit anti DARPP-32 (1:500, sc-11,365, Santa Cruz Biotech); mouse anti D1 (1:500; sc-33,660, Santa Cruz Biotech); goat anti Activin A (1:250, AF338, RnD); mouse anti GFAP-Cy3 conjugated (1:1,000, C9205, Sigma-Aldrich) and rabbit anti Iba1 (1:1,000, Wako, 0719-19,741).

For detection of *Acvr1b* mRNA by *in situ* hybridization, a template for riboprobe generation was obtained by RT-PCR from P1 mouse brain tissue using sense primer, 5'-TTTAAGCTGTTCCTCTGCCTAC-3' and antisense primer, 5'-CCCAAGACTTCCACCTA CATTTC-3' ($T_m = 62$ °C) and subsequently subcloned into PCR II-TOPO-TA cloning vector (Invitrogen). Riboprobes were generated with a BIOT-NTP labeling kit (Roche). For the antisense riboprobe, the template plasmid was linearized with SpeI (New England Biolabs) and transcribed with T7 polymerase (Roche); for the sense riboprobe, the template was linearized with XhoI (New England Biolabs) and transcribed with Sp6 polymerase (Roche). Coronal brain sections of 30 μ m were obtained on a cryostat. Free-floating sections were washed in PBS followed by 5X saline-sodium citrate buffer (SCC) (0.75 M NaCl and 75 mM sodium citrate). The sections were then prehybridized for 2 h at 50°C in SCC buffer containing 50% deionized formamide and 40 μ g/mL salmon DNA (ThermoFisher) and then incubated with sense or antisense probes at 75°C for 10 min, followed by 58°C for 16 h. After washing at 65°C for 2 h, the sections were treated with distilled water to quench endogenous autofluorescence, blocked in 0.5% tyramide signal amplification solution (TSA; TSA biotin kit; PerkinElmer) for 30 min, and stained with Streptavidin-HRP (1:100) at room temperature. After several washes, the sections were incubated in biotin-tyramide (1:100) and reconstituted in amplification diluent (TSA biotin kit; PerkinElmer) for 10 min. After further washes, the sections were stained with Streptavidin-488 (Life Technologies; S32355; 1:1,500 dilution) for 90 min. Sections were then washed and prepared for immunohistochemistry as described above. Control hybridizations with sense riboprobe did not give any signal.

All fluorescence images were captured with a Carl Zeiss LSM 710 confocal microscope using ZEN2012 software (Carl Zeiss). Image analysis was performed using ImageJ and Fiji software packages. Quantification of the number of PLA dots was done in ImageJ using the plugin Synaptic Puncta Analyzer (Ippolito & Eroglu, 2010). The background was subtracted from the image, and the same intensity threshold and range of particle size were used to analyze all images from the same experiment.

Statistical analysis

Statistics analyses were performed using Prism 9 and Graph Plot for macOS (GraphPad Software, LLC.). Data are expressed as average \pm S.E.M. Sample sizes were determined *a priori* as $N = 3\text{--}4$ for *in vitro* studies and $N = 5\text{--}6$ for *in vivo* studies based on previous work from our laboratory and other studies in the field. The *F*-test or Kolmogorov–Smirnov test with Dallal–Wilkinson–Lillie for *P*-value were used to assess data normality and homogeneity variance. Group comparisons were made using the unpaired Student's *t*-test, one-way ANOVA, or two-way ANOVA as appropriate, followed by the Bonferroni or Tukey's multiple comparisons test for normally distributed data. Differences were considered significant when $P < 0.05$. Individual mice and cell culture dishes were randomly assigned to experimental groups to avoid bias. Experimenters were not blind as to the genotype of the cells and mice used in the study.

Data availability

This study does not include data deposited in external repositories.

Expanded View for this article is available online.

Acknowledgements

The authors would like to thank Iina-Lotta Eleonoora Korkala (Karolinska Institute) for help with RT-qPCR; Linda Thors, Emma Wallet, and Jessica Sundell (KMB animal facility, Karolinska Institute) for assistance with animal care; Albert Blanchart Aguado (VirusTech Core, Karolinska Institute) for assistance with the production of shPCBP1-lentivirus; Klas Kullander (Uppsala University, Sweden) for *Gpr101^{Cte}* mice; and all CIBLab members for comments and suggestions. Support for this research was provided by grants to C.F.I. from the Swedish Research Council (2016-01538 and 2020-01923).

Author contributions

Favio A Krapacher: Conceptualization; formal analysis; investigation; methodology. **Diana Fernandez-Suarez:** Investigation; methodology. **Annika Andersson:** Investigation; methodology. **Alvaro Carrier-Ruiz:** Investigation; methodology. **Carlos F Ibanez:** Conceptualization; formal analysis; supervision; funding acquisition; investigation; project administration.

In addition to the CRediT author contributions listed above, the contributions in detail are:

FK performed the majority of experimental work; DFS contributed to experimental design, stereotaxic injection, and *in situ* hybridization; AA performed all mouse genotyping and assisted in cell culture and histology studies; AC-R performed the ALK4 immunohistochemistry; FK and CFI designed the experiments; CFI wrote the paper.

Disclosure and competing interests statement

The authors declare that they have no conflict of interest.

References

- Alibhai IN, Green TA, Potashkin JA, Nestler EJ (2007) Regulation of fosB and DeltafosB mRNA expression: *in vivo* and *in vitro* studies. *Brain Res* 1143: 22–33
- Andersson O, Korach-Andre M, Reissmann E, Ibáñez CF, Bertolino P (2008) Growth/differentiation factor 3 signals through ALK7 and regulates accumulation of adipose tissue and diet-induced obesity. *Proc Natl Acad Sci U S A* 105: 7252–7256
- Andersson O, Reissmann E, Jörnvall H, Ibáñez CF (2006) Synergistic interaction between Gdf1 and Nodal during anterior axis development. *Dev Biol* 293: 370–381
- Bernardi RE, Spanagel R (2014) A two-injection protocol for nicotine sensitization. *Behav Brain Res* 275: 11–14
- Bock R, Shin JH, Kaplan AR, Dobi A, Markey E, Kramer PF, Gremel CM, Christensen CH, Adrover MF, Alvarez VA (2013) Strengthening the accumbal indirect pathway promotes resilience to compulsive cocaine use. *Nat Neurosci* 16: 632–638
- Budi EH, Duan D, Derynck R (2017) Transforming Growth Factor- β Receptors and Smads: Regulatory Complexity and Functional Versatility. *Trends Cell Biol* 27: 658–672
- Burkovetskaya ME, Small R, Guo L, Buch S, Guo M-L (2020) Cocaine self-administration differentially activates microglia in the mouse brain. *Neurosci Lett* 728: 134951
- Chaudhury A, Hussey GS, Ray PS, Jin G, Fox PL, Howe PH (2010) TGF- β -mediated phosphorylation of hnRNP E1 induces EMT via transcript-selective translational induction of Dab2 and ILEI. *Nat Cell Biol* 12: 286–293
- Chen J, Kelz MB, Hope BT, Nakabeppu Y, Nestler EJ (1997) Chronic Fos-related antigens: stable variants of deltaFosB induced in brain by chronic treatments. *J Neurosci* 17: 4933–4941
- Chen J, Nye HE, Kelz MB, Hiroi N, Nakabeppu Y, Hope BT, Nestler EJ (1995) Regulation of delta FosB and FosB-like proteins by electroconvulsive seizure and cocaine treatments. *Mol Pharmacol* 48: 880–889
- Chivero ET, Thangaraj A, Tripathi A, Periyasamy P, Guo M-L, Buch S (2021) NLRP3 Inflammasome Blockade Reduces Cocaine-Induced Microglial Activation and Neuroinflammation. *Mol Neurobiol* 58: 2215–2230
- Clark KH, Wiley CA, Bradberry CW (2013) Psychostimulant abuse and neuroinflammation: emerging evidence of their interconnection. *Neurotox Res* 23: 174–188
- Derynck R, Budi EH (2019) Specificity, versatility, and control of TGF- β family signaling. *Sci Signal* 12: eaav5183
- ten Dijke P, Yamashita H, Ichijo H, Franzén P, Laiho M, Miyazono K, Heldin C-H (1994) Characterization of type I receptors for transforming growth factor- β and activin. *Science* 264: 101–104
- Ersche KD, Dörfinger R (2017) Inflammation and infection in human cocaine addiction. *Curr Opin Behav Sci* 13: 203–209
- Gancarz AM, Wang Z-J, Schroeder GL, Damez-Werno D, Braunscheidel KM, Mueller LE, Humby MS, Caccamise A, Martin JA, Dietz KC et al (2015) Activin receptor signaling regulates cocaine-primed behavioral and morphological plasticity. *Nat Neurosci* 18: 959–961
- Göngrich C, Krapacher FA, Munguba H, Fernandez-Suarez D, Andersson A, Hjerling-Leffler J, Ibáñez CF (2020) ALK4 coordinates extracellular and intrinsic signals to regulate development of cortical somatostatin interneurons. *J Cell Biol* 219: e201905002
- Graybiel AM, Moratalla R, Robertson HA (1990) Amphetamine and cocaine induce drug-specific activation of the c-fos gene in striosome-matrix compartments and limbic subdivisions of the striatum. *Proc Natl Acad Sci U S A* 87: 6912–6916
- Gu Z, Nomura M, Simpson BB, Lei H, Feijen A, van den Eijnden-van RA, Donahoe PK, Li E (1998) The type I activin receptor ActRIB is required for egg cylinder organization and gastrulation in the mouse. *Genes Dev* 12: 844–857
- Hasegawa Y, Mukai H, Asashima M, Hojo Y, Ikeda M, Komatsuzaki Y, Ooishi Y, Kawato S (2014) Acute modulation of synaptic plasticity of pyramidal neurons by activin in adult hippocampus. *Front Neural Circuits* 8: 56

- Hiroi N, Brown JR, Haile CN, Ye H, Greenberg ME, Nestler EJ (1997) FosB mutant mice: loss of chronic cocaine induction of Fos-related proteins and heightened sensitivity to cocaine's psychomotor and rewarding effects. *Proc Natl Acad Sci U S A* 94: 10397–10402
- Hope B, Kosofsky B, Hyman SE, Nestler EJ (1992) Regulation of immediate early gene expression and AP-1 binding in the rat nucleus accumbens by chronic cocaine. *Proc Natl Acad Sci U S A* 89: 5764–5768
- Hope BT, Nye HE, Kelz MB, Self DW, Iadarola MJ, Nakabeppu Y, Duman RS, Nestler EJ (1994) Induction of a long-lasting Ap-1 complex composed of altered fos-like proteins in brain by chronic cocaine and other chronic treatments. *Neuron* 13: 1235–1244
- Hwang CK, Wagley Y, Law P-Y, Wei L-N, Loh HH (2017) Phosphorylation of poly(rC) binding protein 1 (PCBP1) contributes to stabilization of mu opioid receptor (MOR) mRNA via interaction with AU-rich element RNA-binding protein 1 (AUF1) and poly A binding protein (PABP). *Gene* 598: 113–130
- Ippolito DM, Eroglu C (2010) Quantifying synapses: an immunocytochemistry-based assay to quantify synapse number. *J Vis Exp* e2270: <https://doi.org/10.3791/2270>
- Jiang SZ, Sweat S, Dahlke SP, Loane K, Drossel G, Xu W, Tejada HA, Gerfen CR, Eiden LE (2021) Cocaine-dependent acquisition of locomotor sensitization and conditioned place preference requires D1 dopaminergic signaling through a cyclic AMP, NCS-Rapgef2, ERK, and Egr-1/Zif268 pathway. *J Neurosci* 41: 711–725
- Kauer JA, Malenka RC (2007) Synaptic plasticity and addiction. *Nat Rev Neurosci* 8: 844–858
- Kelz MB, Chen J, Carlezon WA, Whisler K, Gilden L, Beckmann AM, Steffen C, Zhang YJ, Marotti L, Self DW et al (1999) Expression of the transcription factor deltaFosB in the brain controls sensitivity to cocaine. *Nature* 401: 272–276
- Kravitz AV, Tye LD, Kreitzer AC (2012) Distinct roles for direct and indirect pathway striatal neurons in reinforcement. *Nat Neurosci* 15: 816–818
- Liao K, Guo M, Niu F, Yang L, Callen SE, Buch S (2016) Cocaine-mediated induction of microglial activation involves the ER stress-TLR2 axis. *J Neuroinflammation* 13: 33
- Lüscher C (2016) The emergence of a circuit model for addiction. *Annu Rev Neurosci* 39: 257–276
- Marinescu V, Loomis PA, Ehmann S, Beales M, Potashkin JA (2007) Regulation of retention of FosB intron 4 by PTB. *PLoS One* 2: e828
- Massagué J (2012) TGF β signalling in context. *Nat Rev Mol Cell Biol* 13: 616–630
- Meng Q, Rayala SK, Gururaj AE, Talukder AH, O'Malley BW, Kumar R (2007) Signaling-dependent and coordinated regulation of transcription, splicing, and translation resides in a single coregulator, PCBP1. *Proc Natl Acad Sci U S A* 104: 5866–5871
- Moustakas A, Heldin C-H (2009) The regulation of TGFbeta signal transduction. *Development* 136: 3699–3714
- Mumberg D, Lucibello FC, Schuermann M, Müller R (1991) Alternative splicing of fosB transcripts results in differentially expressed mRNAs encoding functionally antagonistic proteins. *Genes Dev* 5: 1212–1223
- Nakabeppu Y, Nathans D (1991) A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity. *Cell* 64: 751–759
- Nestler EJ (2008) Transcriptional mechanisms of addiction: role of DeltaFosB. *Philos Trans R Soc Lond B Biol Sci* 363: 3245–3255
- Nestler EJ (2013) Δ FosB: a molecular switch for reward. *J Drug Alcohol Res* 2: 1–11
- Nestler EJ, Barrot M, Self DW (2001) DeltaFosB: a sustained molecular switch for addiction. *Proc Natl Acad Sci U S A* 98: 11042–11046
- Peakman M-C, Colby C, Perrotti LI, Tekumalla P, Carle T, Ulery P, Chao J, Duman C, Steffen C, Monteggia L et al (2003) Inducible, brain region-specific expression of a dominant negative mutant of c-Jun in transgenic mice decreases sensitivity to cocaine. *Brain Res* 970: 73–86
- Penrod RD, Kourrich S, Kearney E, Thomas MJ, Lanier LM (2011) An embryonic culture system for the investigation of striatal medium spiny neuron dendritic spine development and plasticity. *J Neurosci Methods* 200: 1–13
- Penrod RD, Thomsen M, Taniguchi M, Guo Y, Cowan CW, Smith LN (2020) The activity-regulated cytoskeleton-associated protein, Arc/Arg3.1, influences mouse cocaine self-administration. *Pharmacol Biochem Behav* 188: 172818
- Reinius B, Blunder M, Brett FM, Eriksson A, Patra K, Jonsson J, Jazin E, Kullander K (2015) Conditional targeting of medium spiny neurons in the striatal matrix. *Front Behav Neurosci* 9: 71
- Renthal W, Kumar A, Xiao G, Wilkinson M, Covington HEI, Maze I, Sikder D, Robison AJ, LaPlant Q, Dietz DM et al (2009) Genome-wide analysis of chromatin regulation by cocaine reveals a role for sirtuins. *Neuron* 62: 335–348
- Robison AJ, Nestler EJ (2022) Δ FOSB: a potentially druggable master orchestrator of activity-dependent gene expression. *ACS Chem Neurosci* 13: 296–307
- Roussis IM, Guille M, Myers FA, Scarlett GP (2016) RNA whole-mount *in situ* hybridisation proximity ligation assay (riSH-PLA), an assay for detecting rna-protein complexes in intact cells. *PLoS One* 11: e0147967
- Ruffle JK (2014) Molecular neurobiology of addiction: what's all the (Δ)Fos about? *Am J Drug Alcohol Abuse* 40: 428–437
- Schmierer B, Hill CS (2007) TGFbeta-SMAD signal transduction: molecular specificity and functional flexibility. *Nat Rev Mol Cell Biol* 8: 970–982
- Su X, Huang L, Xiao D, Qu Y, Mu D (2018) Research progress on the role and mechanism of action of activin A in brain injury. *Front Neurosci* 12: 697
- Teague CD, Nestler EJ (2021) Key transcription factors mediating cocaine-induced plasticity in the nucleus accumbens. *Mol Psychiatry* 27: 687–709
- Tolu S, Avale ME, Nakatani H, Pons S, Parnaudeau S, Tronche F, Vogt A, Monyer H, Vogel R, de Chaumont F et al (2010) A versatile system for the neuronal subtype specific expression of lentiviral vectors. *FASEB J* 24: 723–730
- Tripathi V, Sixt KM, Gao S, Xu X, Huang J, Weigert R, Zhou M, Zhang YE (2016) Direct regulation of alternative splicing by SMAD3 through PCBP1 is essential to the tumor-promoting role of TGF- β . *Mol Cell* 64: 1010
- Tsuchida K, Nakatani M, Yamakawa N, Hashimoto O, Hasegawa Y, Sugino H (2004) Activin isoforms signal through type I receptor serine/threonine kinase ALK7. *Mol Cell Endocrinol* 220: 59–65
- Valjent E, Aubier B, Corbillé A-G, Brami-Cherrier K, Caboche J, Topilko P, Girault J-A, Hervé D (2006) Plasticity-associated gene Krox24/Zif268 is required for long-lasting behavioral effects of cocaine. *J Neurosci* 26: 4956–4960
- Valjent E, Bertran-Gonzalez J, Aubier B, Greengard P, Hervé D, Girault J-A (2010) Mechanisms of locomotor sensitization to drugs of abuse in a two-injection protocol. *Neuropsychopharmacology* 35: 401–415
- Verschueren K, Dewulf N, Goumans MJ, Lonnoy O, Feijen A, Grimsby S, Vandi Spiegle K, ten Dijke P, Morén A et al (1995) Expression of type I and type IB receptors for activin in midgestation mouse embryos suggests distinct functions in organogenesis. *Mech Dev* 52: 109–123
- Volkow ND, Morales M (2015) The brain on drugs: from reward to addiction. *Cell* 162: 712–725

- Wang Z-J, Martin JA, Gancarz AM, Adank DN, Sim FJ, Dietz DM (2017) Activin A is increased in the nucleus accumbens following a cocaine binge. *Sci Rep* 7: 43658
- Xu Q, Tam M, Anderson SA (2008) Fate mapping Nkx2.1-lineage cells in the mouse telencephalon. *J Comp Neurol* 506: 16–29
- Young ST, Porrino LJ, Iadarola MJ (1991) Cocaine induces striatal c-fos-immunoreactive proteins via dopaminergic D1 receptors. *Proc Natl Acad Sci U S A* 88: 1291–1295
- Zachariou V, Bolanos CA, Selley DE, Theobald D, Cassidy MP, Kelz MB, Shaw-Lutchman T, Berton O, Sim-Selley LJ, Dileone RJ et al (2006) An essential role for DeltaFosB in the nucleus accumbens in morphine action. *Nature* 9: 205–211

- Zhang W, Xie M, Shu M-D, Steitz JA, DiMaio D (2016) A proximity-dependent assay for specific RNA-protein interactions in intact cells. *RNA* 22: 1785–1792



License: This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




Figure S1. Control experiments in cultured MSNs exposed to single or consecutive stimulations

(A) Schematic of the single stimulation protocol used for experiments shown in panels B to H. MSNs were extracted at embryonic day 16.5 (E16.5) and cultured for 7 days (7div) before 2 hours (2hr) treatment with either 1 μ M SKF81297 (SKF), 100ng/ml activin A (Act) or their combination (Act+SKF), after which cultures were fixed for imaging or collected for RNA extraction. (B-F) Real-time quantitative PCR (RT-qPCR) analysis of Δ FosB (B), FosB (C), c-Fos (D), ARC (E) and EGR1 (F) mRNAs in cultured MSNs exposed to a single stimulation with activin A (Act), SKF81297 (SKF) or their combination as indicated. Results are expressed as average \pm SEM of fold increase over vehicle. *, $p < 0.05$; N=5; one-way ANOVA followed with Tukey's multiple comparisons test. (G) Representative immunofluorescence staining of Δ FosB (green) in cultured MSNs exposed to a single stimulation with activin A (Act), SKF81297 (SKF) or their combination as indicated. Counterstaining with MAP2 and DAPI is shown in the lower row. Scale bar, 25 μ m. (H) Quantification of Δ FosB+ nuclei in cultured MSNs after the indicated single stimulation treatments. Results are expressed as average \pm SEM of % of MAP2+ cells displaying Δ FosB in the nucleus. No significant differences were found by one-way ANOVA ($p \geq 0.05$; N=6). (I) RT-qPCR analysis of *Acvr1b* mRNA (encoding ALK4) in cultured MSNs

exposed to consecutive stimulation as indicated (same treatment in both treatment days). Results are expressed as average \pm SEM of fold change over vehicle. No significant differences were found by one-way ANOVA followed with Tukey's multiple comparisons test ($p \geq 0.05$; N=4).

(J) RT-qPCR analysis of Δ FosB mRNA in cultured MSNs exposed to consecutive stimulation (same treatment in both treatment days) with activin A (Act), SKF81297 (SKF), SB431542 (SB) or PKA inhibitor (PKI) as indicated. Results are expressed as average \pm SEM of fold change over vehicle. **, $p < 0.01$; ***, $p < 0.001$; N=4; one-way ANOVA followed with Tukey's multiple comparisons test.

(K) Representative immunofluorescence staining of Δ FosB (green) in cultured MSNs exposed to consecutive stimulation (same treatment in both treatment days) with activin A (Act), SKF81297 (SKF) SB431542, (SB) or PKA inhibitor (PKI) as indicated. Counterstaining with MAP2 and DAPI is shown in the lower row. Scale bar, 25 μ m.

(L) Quantification of Δ FosB+ nuclei in cultured MSNs after the indicated treatments (same treatment in both treatment days). Results are expressed as average \pm SEM of % of MAP2+ cells displaying Δ FosB in the nucleus. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; N=6; one-way ANOVA followed with Tukey's multiple comparisons test.




Figure S2. Expression of Acvr1b mRNA and ALK4 protein in nucleus accumbens of wild type and *Gad67^{Cre};Alk4^{fl/fl}* mutant mice

(A) Representative *in situ* hybridization of Acvr1b mRNA (red) and immunofluorescence of DARPP-32 (green) and D1 dopamine receptor (grey) in NAc and striatum of wild type mice. Scale bar, 25 μ M.

(B, C) Quantification of cells expressing Acvr1b mRNA (Acvr1b⁺) in NAc and striatum of wild type mice. Results are expressed as average \pm SEM of % of DARPP-32⁺ (B) or D1⁺ (C) cells displaying signal for Acvr1b mRNA. N=4.

(D) Representative *in situ* hybridization of Acvr1b mRNA (red) and immunofluorescence of DARPP-32 (green) in N. accumbens and striatum of control (*Gad67^{Cre}*) and mutant (*Gad67^{Cre};Alk4^{fl/fl}*) mice. Scale bar, 25 μ M.

(E, F) RT-qPCR analysis of Acvr1b mRNA in N. accumbens

(E) and primary motor cortex (F) of control (*Gad67^{Cre}*) and mutant (*Gad67^{Cre};Alk4^{fl/fl}*) mice. Results are expressed as average \pm SEM of fold change over control mice. **, p<0.01; N=6; Student's t-test. No significant differences were found in cortex (p \geq 0.05).

(G) Representative immunofluorescence staining of ALK4 (red) and DARPP-32 (green) in N. accumbens of control (*Gad67^{Cre}*) and mutant (*Gad67^{Cre};Alk4^{fl/fl}*) mice. Scale bar, 25 μ M.

(H) Representative immunofluorescence staining of ALK4 in primary motor cortex of control (*Gad67^{Cre}*) and mutant (*Gad67^{Cre};Alk4^{fl/fl}*) mice. Scale bar, 50 μ M.




Figure S3. RNA-PLA between PCBP1 or Smad3 with FosB mRNA: regulated intron 4 and control probes

(A) Representative images of RNA-PLA (green) between PCBP1 (left) or Smad3 (right) and sequences in regulated intron 4 (prl4) of FosB mRNA, respectively, in HeLa cells exposed to consecutive stimulation (same treatment in both treatment days) with activin A (Act), forskolin (Fsk), SB431542 (SB), or PKA inhibitor (PKI) as indicated. Counterstaining with phalloidin and DAPI is shown in the second and fourth columns. Scale bars, 20 μ M.

(B, C) Quantification of RNA-PLA between PCBP1 (B) or Smad3 (C) and prl4 in HeLa cells after the indicated treatments. Results are expressed as average \pm SEM of number of RNA-PLA dots per cell. Only nuclear dots were quantified in this analysis. *, $p < 0.05$; ***, $p < 0.001$; N=4; one-way ANOVA followed with Tukey's multiple comparisons test.

(D, E) Representative images of control probes for RNA-PLA (red) between PCBP1 (D) or Smad3 (E) and sequences in exon 3 (pE3), exon 4a/b junction (pE4a/b), exon 4 (pE4) sense and regulated intron 4 (prl4) sense of FosB mRNA, respectively, in HeLa cells exposed to consecutive stimulation (same treatment in both treatment days) with Act+SKF or vehicle, as indicated. Counterstaining with phalloidin and DAPI is shown for each panel. Scale bars, 20 μ M.

(F) Representative images of control antibody (against Sirt1) for RNA-PLA (red) and sequences in exon 4 (pE4) and regulated intron 4 (prl4) of FosB mRNA, respectively, in HeLa cells exposed to consecutive stimulation as indicated. Counterstaining with phalloidin and DAPI is shown for each panel. Scale bars, 20 μ M.




Figure S4. Expression of Acvr1b mRNA in nucleus accumbens of wild type mice, *Nkx2.1Cre;Alk4fl/fl* and *Gpr101Cre;Alk4fl/fl* mutant mice and after AAV2-Cre injection

(A) RT-qPCR analysis of *Acvr1b* mRNA in N. accumbens of control (*Nkx2.1^{Cre}*) and mutant (*Nkx2.1^{Cre};Alk4^{fl/fl}*) mice. Results are expressed as average \pm SEM of fold change over control mice. No significant differences were found between genotypes ($p \geq 0.05$). N=6; Student's t-test.

(B, C) RT-qPCR analysis of *Acvr1b* mRNA in N. accumbens (B) and primary motor cortex (C) of control (*Gpr101^{Cre}*) and mutant (*Gpr101^{Cre}; Alk4^{fl/fl}*) mice. Results are expressed as average \pm SEM of fold change over control mice. ***, $p < 0.001$; N=6; Student's t-test. No significant differences were found in cortex ($p \geq 0.05$).

(D) Representative in situ hybridization of *Acvr1b* mRNA (red) and immunofluorescence of DARPP-32 (green) in N. accumbens and striatum of control (*Gpr101^{Cre}*) and mutant

(*Gpr101^{Cre}; Alk4^{fl/fl}*) mice. Scale bar, 25 μ M.

(E) Representative fluorescence of AAV2-GFP (green) viral injection site in NAc shell. NAc core and dorsal striatum (dStr) are indicated. Lower magnification shown in left panel, scale bar 400 μ M. Central and right panels show higher magnification (scale bar, 100 μ M), with immunofluorescence staining for DARPP-32 (red) to the right.

(F, G) RT-qPCR analysis of *Acvr1b* mRNA in N. accumbens (F) and primary motor cortex (G) of *Alk4^{fl/fl}* mice that received control (AAV2-GFP) or Cre-expressing (AAV2-Cre/GFP) viruses. Results are expressed as average \pm SEM of fold change over AAV2-GFP. **, $p < 0.01$; N=6; Student's t-test. No significant differences were found in cortex ($p \geq 0.05$).




Figure S5. Activin A is induced in NAc microglia after cocaine TIPS

(A, B) RT-qPCR analysis of Inhba (A), encoding activin A and Gdf-1 (B) mRNAs in primary motor cortex (Ctx), hippocampus (Hipp), NAc and mid brain (MdBrain) of wild type mice. Results are expressed as average \pm SEM of fold change over Ctx.

(C) Representative immunofluorescence staining of DARPP-32 (red, first row), GFAP (red, second row) and Iba1 (red, third row), and activin A (green) in coronal sections through the NAc of wild type mice. Activin A immunoreactivity was found to colocalize with all three markers. Scale bars, 20 μ M.

(D) Representative immunofluorescence staining of Iba1 (red) and activin A (green) showing the overlap between the two markers at high magnification in individual microglia cells in NAc and striatum of wild type mice. Scale bar, 10 μ M

(E, F) RT-qPCR analysis of Inhba in NAc of wild type mice subjected to cocaine TIPS 1 hour after a single injection of saline (Sal) or cocaine (Coc) (E) or 1 hour after the second injection (F) the 4 groups of mice of TIPS, as indicated. Results are expressed as

average \pm SEM of fold increase over the saline group (Sal or Sal/Sal, respectively). **, $p < 0.01$; $N = 4$; one-way ANOVA followed with Tukey's multiple comparisons test. No significant differences were found in after the first injection (E) ($p \geq 0.05$).

(G, I) Representative immunofluorescence staining of Iba1 (G, red) or DARPP-32 (I, red) and activin A (green) in coronal sections through the NAc of wild type mice subjected to cocaine TIPS 1 hour after the second injection. Only the Sal/Sal and Coc/Coc (1st injection/2nd injection) groups are shown. Scale bars, 20 μ M.

(H, J) Quantification of cells immunoreactive for activin A (ActA+) in NAc of wild type mice subjected to cocaine TIPS 1 hour after the second injection. Only the Sal/Sal and Coc/Coc groups are shown. Results are expressed as average \pm SEM of % of ActA+ cells among the Iba1+ (H) or DARPP-32+ (J) cell populations. *, $p < 0.05$; $N = 4$; Unpaired t test. No significant differences were found in % of ActA+ cells among the DARPP-32+ population (J) ($p \geq 0.05$).




Figure S6. ALK4 is required in NAc MSNs of adult mice for sustained induction of Δ FosB mRNA during cocaine sensitization

RT-qPCR analysis of FosB (A, D), c-Fos (B, E) and Δ FosB (C, F) mRNAs in NAc of *Gpr101^{Cre};Alk4^{fl/fl}* mutant mice 1 hour (A-C) or 24 hour (D-F) after the second saline (Sal) or cocaine (Coc) injection of TIPS. Results are expressed as average \pm SEM of fold increase over the saline group (Sal/Sal) of control mice. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $N = 5$; two-way ANOVA followed with Tukey's multiple comparisons test.




Figure S7. ALK4 is required in NAC MSNs of adult mice for sustained induction of Δ FosB targets ARC and EGR1 mRNAs during cocaine sensitization

(A, B) RT-qPCR analysis of ARC (A) and EGR1 (B) mRNAs in NAC of *Gad67^{Cre};Alk4^{fl/fl}* mutant mice 24 hour after the second saline (Sal) or cocaine (Coc) injection of TIPS using conditions identical to those of Figure 6. Results are expressed as average \pm SEM of fold increase over the saline group (Sal/Sal) of control mice (*Gad67^{Cre}*). ***, $p < 0.001$; $N = 5$; two-way ANOVA followed with Tukey's multiple comparisons test.

(C, D) RT-qPCR analysis of ARC (A) and EGR1 (B) mRNAs in NAC of *Gpr101^{Cre};Alk4^{fl/fl}* mutant mice 24 hour after the second saline (Sal) or cocaine (Coc) injection of TIPS using conditions identical to those of Figure 6. Results are expressed as average \pm SEM of fold increase over the saline group (Sal/Sal) of control mice (*Gpr101^{Cre}*). **, $p < 0.01$; ***, $p < 0.001$; $N = 5$; two-way ANOVA followed with Tukey's multiple comparisons test.




Figure S8. Schematic of the pathway and molecular relationships implicated by the results of this study

PKA activation downstream of D1 signaling phosphorylates PCBP1. ALK4 activation following activin A binding phosphorylates Smad3. Phosphorylated PCBP1 and Smad3 form a complex that translocates to the nucleus. In the nucleus, the P-PCBP1/P-Smad3 complex enhances splicing of the regulated intron (r14) of *FosB* mRNA, possibly, by regulating the function of the U2/U2AF small nuclear ribonucleoprotein complex (dashed arrow). Removal of r14 brings the cryptic stop codon (red line) in-frame, resulting in the much shorter Δ FosB protein.